




ALGEBRAIC COMBINATORICS

Yu-Cheng Chiu & Eric Marberg
Expanding K -theoretic Schur Q -functions
Volume 6, issue 6 (2023), p. 1419-1445.
<https://doi.org/10.5802/alco.312>

© The author(s), 2023.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION (CC-BY) 4.0 LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*Algebraic Combinatorics is published by The Combinatorics Consortium
and is a member of the Centre Mersenne for Open Scientific Publishing*
www.tccpublishing.org www.centre-mersenne.org
e-ISSN: 2589-5486





Expanding K -theoretic Schur Q -functions

Yu-Cheng Chiu & Eric Marberg

ABSTRACT We derive several identities involving Ikeda and Naruse’s K -theoretic Schur P - and Q -functions. Our main result is a formula conjectured by Lewis and the second author which expands each K -theoretic Schur Q -function in terms of K -theoretic Schur P -functions. This formula extends to some more general identities relating the skew and dual versions of both power series. We also prove a shifted version of Yeliussizov’s skew Cauchy identity for symmetric Grothendieck polynomials. Finally, we discuss some conjectural formulas for the dual K -theoretic Schur P - and Q -functions of Nakagawa and Naruse. We show that one such formula would imply a basis property expected of the K -theoretic Schur Q -functions.

1. INTRODUCTION

This article proves some identities relating the K -theoretic Schur P - and Q -functions introduced by Ikeda and Naruse in [8]. To motivate the definition of these power series and to frame our main results, we start by reviewing some classical background material on generating functions for shifted tableaux.

Let $\lambda = (\lambda_1 > \lambda_2 > \cdots > 0)$ be a *strict partition*, that is, a strictly decreasing sequence of positive integers. The *shifted diagram* of λ is the set of pairs $\text{SD}_\lambda := \{(i, j) \in \mathbb{Z} \times \mathbb{Z} : 0 < i \leq j < i + \lambda_i\}$. We usually refer to the elements of this set as “positions” or “boxes.”

A *shifted tableau* of shape λ is a filling of SD_λ by positive half-integers. For any $i \in \mathbb{Z}$ let $i' := i - \frac{1}{2}$. Then one may think of the entries of a shifted tableau as consisting of positive integers i and primed numbers i' . A shifted tableau is *semistandard* if the following conditions hold:

- (S1) The entries in each row and column are weakly increasing.
- (S2) No unprimed number i occurs more than once in a given column.
- (S3) No primed number i' occurs more than once in a given row.

Let $\text{ShYT}_Q(\lambda)$ denote the set of semistandard shifted tableaux of shape λ . Define $\text{ShYT}_P(\lambda) \subseteq \text{ShYT}_Q(\lambda)$ to be the subset of tableaux also satisfying:


- (S4) No primed number occurs in any diagonal position $(j, j) \in \text{SD}_\lambda$.

Manuscript received 22nd December 2021, revised 31st October 2022, accepted 11th April 2023.

KEYWORDS. K -theoretic Schur P - and Q -functions, Cauchy identities, shifted tableaux, set-valued tableaux, plane partitions.

ACKNOWLEDGEMENTS. The authors were partially supported by grants ECS 26305218 and GRF 16306120 from the Hong Kong Research Grants Council.

We refer to elements of $\text{ShYT}_P(\lambda)$ and $\text{ShYT}_Q(\lambda)$ as *P-shifted* and *Q-shifted tableaux*, respectively. We draw shifted tableaux in French notation:



The *weight* of a shifted tableau T is the monomial $x^T := \prod_{i \geq 1} x_i^{m_i}$ where m_i is the number of times that i or i' appears in T . For example, we have $x^T = x_1 x_2^2 x_3^3 x_4$ for both of the shifted tableaux shown as examples above. The *Schur P- and Q-functions* indexed by λ are the power series

$$(1.1) \quad P_\lambda := \sum_{T \in \text{ShYT}_P(\lambda)} x^T \quad \text{and} \quad Q_\lambda := \sum_{T \in \text{ShYT}_Q(\lambda)} x^T.$$


Any way of toggling the primes in the diagonal entries of a *Q*-shifted tableau results in another *Q*-shifted tableau of the same weight, so it is clear that

$$(1.2) \quad Q_\lambda = 2^{\ell(\lambda)} P_\lambda \quad \text{where} \quad \ell(\lambda) := |\{i : \lambda_i > 0\}| = |\{i : (i, i) \in \text{SD}_\lambda\}|.$$

It is well-known that P_λ and Q_λ are symmetric functions of bounded degree. They were first defined in work of Schur on the projective representations of the symmetric group but have since appeared in various other contexts.

We are interested in generalizations of P_λ and Q_λ that are similar generating functions for set-valued shifted tableaux. A *set-valued shifted tableau* of shape λ is a filling of SD_λ by nonempty finite subsets of $\{\frac{1}{2}i : 0 < i \in \mathbb{Z}\} = \{1' < 1 < 2' < 2 < \dots\}$. We consider a sequence of such subsets S_1, S_2, S_3, \dots to be *weakly increasing* if $\max(S_i) \leq \min(S_{i+1})$ for all i . With this convention, we may define a set-valued shifted tableau to be *semistandard* if it satisfies the same conditions (S1)–(S3) as above.

We write $\text{SetShYT}_Q(\lambda)$ for the set of all semistandard set-valued shifted tableaux of shape λ , and $\text{SetShYT}_P(\lambda)$ for the subset of such tableaux also satisfying (S4). We refer to elements of $\text{SetShYT}_P(\lambda)$ and $\text{SetShYT}_Q(\lambda)$ as *set-valued P-shifted* and *set-valued Q-shifted tableaux*, respectively:



The *weight* x^T of a set-valued shifted tableau T is defined in the same way as in the non-set-valued case; for both tableaux in the preceding example one has $x^T = x_1 x_2^3 x_3^4 x_4 x_5$. Write T_{ij} for the entry of a set-valued shifted tableau in position (i, j) and define $|T| := \sum_{(i,j) \in \text{SD}_\lambda} |T_{ij}|$ and $|\lambda| := |\text{SD}_\lambda|$. Then $|T| - |\lambda|$ is the difference between the degree of x^T and the size of SD_λ . Finally, let β be a variable that commutes with each x_i . The *K-theoretic Schur P- and Q-functions* indexed by λ are the power series

in $\mathbb{Z}[\beta][[x_1, x_2, \dots]]$ given by

$$(1.3) \quad \begin{aligned} GP_\lambda &:= \sum_{T \in \text{SetShYT}_P(\lambda)} \beta^{|T| - |\lambda|} x^T, \\ GQ_\lambda &:= \sum_{T \in \text{SetShYT}_Q(\lambda)} \beta^{|T| - |\lambda|} x^T. \end{aligned}$$

We recover P_λ from GP_λ and Q_λ from GQ_λ by setting $\beta = 0$. Both GP_λ and GQ_λ are symmetric in the x_i variables [8, §3.4] and homogeneous of degree $|\lambda|$ if we set $\deg(\beta) = -1$ and $\deg(x_i) = 1$.

Ikedo and Naruse introduced these functions in [8] for applications in K -theory. Specializations of GP_λ and GQ_λ represent the structure sheaves of Schubert varieties in the K -theory of the maximal isotropic Grassmannians of orthogonal and symplectic types [8, Cor. 8.1]. More precisely, the GP - and GQ -functions represent Schubert classes in connective K -theory, so can be turned into cohomology classes or elements of the Grothendieck ring of vector bundles on setting $\beta = 0$ and $\beta = 1$, respectively. The GP - and GQ -functions are also “stable limits” of connective K -theory classes of orbit closures for symplectic and orthogonal groups acting on the type A flag variety [14, 15]. For more results about these functions and various extensions, see [16, 17, 18].

Our first main result is a K -theoretic analogue of equation (1.2). The relevant identity is subtler than in the classical case, and was predicted as [11, Conj. 5.15]. It expresses each K -theoretic Schur Q -function as a finite linear combination of K -theoretic Schur P -functions with integer coefficients.

If $\lambda = (\lambda_1 > \lambda_2 > \dots > 0)$ and $\mu = (\mu_1 > \mu_2 > \dots > 0)$ are strict partitions with $\mu_i \leq \lambda_i$ for all i then we write $\mu \subseteq \lambda$ and define $\text{SD}_{\lambda/\mu} := \text{SD}_\lambda \setminus \text{SD}_\mu$ and $|\lambda/\mu| := |\text{SD}_{\lambda/\mu}|$. We also let $\text{cols}(\lambda/\mu) := |\{j : (i, j) \in \text{SD}_{\lambda/\mu} \text{ for some } i\}|$ denote the number of distinct columns occupied by the positions in $\text{SD}_{\lambda/\mu}$.

THEOREM 1.1. *If μ is a strict partition with $\ell(\mu)$ parts then*

$$(1.4) \quad GQ_\mu = 2^{\ell(\mu)} \sum_{\lambda} (-1)^{\text{cols}(\lambda/\mu)} (-\beta/2)^{|\lambda/\mu|} GP_\lambda$$

where the sum is over strict partitions $\lambda \supseteq \mu$ with $\ell(\lambda) = \ell(\mu)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip, that is, a subset with at most one position in each row.

For example, it holds that $GQ_{(3,2)} = 4GP_{(3,2)} + 2\beta GP_{(4,2)} - \beta^2 GP_{(4,3)}$ and $GQ_{(n)} = 2GP_{(n)} + \beta GP_{(n+1)}$ for all integers $n > 0$. This formula applies even when $\mu = \emptyset$ is the empty partition, as then the sum has only one term indexed by $\lambda = \emptyset$, giving $GQ_\emptyset = GP_\emptyset = 1$. We prove Theorem 1.1 in Section 3.

The additional complexity in (1.4) compared to (1.2) is related to the fact that in a set-valued Q -shifted tableau, a diagonal entry may contain both i and i' . When this happens there is no simple way to remove all primes from the diagonal without changing the relevant weight.

As a corollary, we may classify when only positive coefficients appear in the GP -expansion of GQ_μ . Let $\mathbb{N} := \{0, 1, 2, \dots\}$.

COROLLARY 1.2. *If μ is a strict partition then GQ_μ is an $\mathbb{N}[\beta]$ -linear combination of GP -functions if and only if all distinct parts of μ differ by at least two.*

Proof. It suffices by Theorem 1.1 to observe that there exists a strict partition $\lambda \supseteq \mu$ with $\ell(\lambda) = \ell(\mu)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip and $\text{cols}(\lambda/\mu) \not\equiv |\lambda/\mu| \pmod{2}$ if and only if $\mu_i - \mu_{i+1} = 1$ for some $i \in [\ell(\mu) - 1]$. \square

We also prove a few more results. Theorem 1.1 has some enumerative consequences which we discuss in Section 4.

The GP - and GQ -functions have skew versions $GP_{\lambda/\mu}$ and $GQ_{\lambda/\mu}$, which are generating functions for set-valued tableaux of shifted skew shapes. In Section 5 we derive an extension of Theorem 1.1 for these power series, along with some other related identities.

There are also dual power series gp_{λ} and gq_{λ} defined by Nakagawa and Naruse [16] from GP_{λ} and GQ_{λ} via a Cauchy identity. Section 6 contains some further results about these functions, including a dual form of Theorem 1.1 (see Corollary 6.2) and a skew Cauchy identity (see Theorem 6.11).

In Section 7, we recall a conjectural formula for gp_{λ} and gq_{λ} from [16]. We then explain a new conjectural formula for the related functions $jp_{\lambda} := \omega(gp_{\lambda})$ and $jq_{\lambda} := \omega(gq_{\lambda})$ obtained by applying the algebra automorphism ω that sends $s_{\lambda} \mapsto s_{\lambda^{\tau}}$. We show that these new conjectures would imply a conjecture of Ikeda and Naruse about the GQ -functions forming a $\mathbb{Z}[\beta]$ -basis for a ring.

2. PRELIMINARIES

Fix a positive integer n and continue to let β, x_1, x_2, \dots be commuting variables. For any $f \in \mathbb{Z}[\beta][[x_1, x_2, \dots]]$ let $f(x_1, x_2, \dots, x_n) \in \mathbb{Z}[\beta][x_1, x_2, \dots, x_n]$ be the polynomial obtained by setting $x_{n+1} = x_{n+2} = \dots = 0$. Also define

$$(2.1) \quad x \oplus y := x + y + \beta xy \quad \text{and} \quad x \ominus y := \frac{x-y}{1+\beta y}.$$

If $\lambda = (\lambda_1, \lambda_2, \dots)$ is a finite sequence of integers then let $x^{\lambda} := \prod_i x_i^{\lambda_i}$.

Ikeda and Naruse use the following formulas as their definition of the K -theoretic Schur P - and Q -functions [8, Def. 2.1]. They derive the set-valued tableau generating functions given in the introduction as [8, Thm. 9.1].

THEOREM 2.1 (See [8]). *If λ is a strict partition with $r := \ell(\lambda) \leq n$ then*

$$GP_{\lambda}(x_1, x_2, \dots, x_n) = \frac{1}{(n-r)!} \sum_{w \in S_n} w \left(x^{\lambda} \prod_{i=1}^r \prod_{j=i+1}^n \frac{x_i \oplus x_j}{x_i \ominus x_j} \right),$$

$$GQ_{\lambda}(x_1, x_2, \dots, x_n) = \frac{1}{(n-r)!} \sum_{w \in S_n} w \left(x^{\lambda} \prod_{i=1}^r (2 + \beta x_i) \prod_{j=i+1}^n \frac{x_i \oplus x_j}{x_i \ominus x_j} \right),$$

where $w \in S_n$ acts on rational functions by permuting the x_i variables while fixing β .

For classical background on the Schur P and Q -functions, see [12, §III.8], [22, §5-§9], or the appendix in [23]. As λ ranges over all strict partitions the functions P_{λ} (respectively, Q_{λ}) are a \mathbb{Z} -basis for a subring of $\mathbb{Z}[[x_1, x_2, \dots]]$. The same is true of the polynomials $P_{\lambda}(x_1, x_2, \dots, x_n)$ (respectively, $Q_{\lambda}(x_1, x_2, \dots, x_n)$) if λ ranges over strict partitions with $\ell(\lambda) \leq n$ [12, §III.8].

Since $P_{\lambda} = GP_{\lambda}|_{\beta=0}$ and $Q_{\lambda} = GQ_{\lambda}|_{\beta=0}$ it follows that the GP_{λ} 's (respectively, the GQ_{λ} 's) are linearly independent over $\mathbb{Z}[\beta]$, as are the polynomials $GP_{\lambda}(x_1, x_2, \dots, x_n)$ (respectively, $GQ_{\lambda}(x_1, x_2, \dots, x_n)$) as λ ranges over all strict partitions with at most n parts.

In fact, Ikeda and Naruse show that the sets $\{GP_{\lambda}(x_1, x_2, \dots, x_n) : \ell(\lambda) \leq n\}$ and $\{GQ_{\lambda}(x_1, x_2, \dots, x_n) : \ell(\lambda) \leq n\}$ are both $\mathbb{Z}[\beta]$ -bases for subrings of $\mathbb{Z}[\beta][x_1, x_2, \dots, x_n]$ [8, Thm. 3.1 and Prop. 3.2]. An analogous basis property is known to hold for the set of all formal power series GP_{λ} 's and is expected to hold for the GQ_{λ} 's; see the discussion before Corollary 7.9.

The GP - and GQ -power series are generalizations of Ivanov's factorial P - and Q -functions [9], which have another generalization studied by Okada in [19]. Comparing

Theorem 2.1 and [19, Lem. 2.4] suggests a common generalization of these functions which might be interesting to consider in future work.

3. EXPANSIONS

We prove Theorem 1.1 in this section. For $m \in \mathbb{N}$ let $[m] := \{1, 2, \dots, m\}$. Fix an integer $n > 0$. Given a nonzero vector $\lambda \in \mathbb{N}^n$, we define rational functions

$$(3.1) \quad \begin{aligned} A_\lambda &:= \frac{1}{r!} \sum_{w \in S_r} w \left(x^\lambda \prod_{i=1}^r \prod_{j=i+1}^n \frac{x_i \oplus x_j}{x_i \ominus x_j} \right), \\ B_\lambda &:= \frac{1}{r!} \sum_{w \in S_r} w \left(x^\lambda \prod_{i=1}^r (2 + \beta x_i) \prod_{j=i+1}^n \frac{x_i \oplus x_j}{x_i \ominus x_j} \right), \end{aligned}$$

where $r := \max\{i \in [n] : \lambda_i \neq 0\}$. Note the implicit dependence on n in these formulas. For convenience we also set $A_0 = B_0 = 1$.

LEMMA 3.1. *Let λ be a strict partition with $r := \ell(\lambda) \leq n$. Fix $m \in [r]$. Let*

$$\mu := (\lambda_1 > \lambda_2 > \dots > \lambda_m) \quad \text{and} \quad \nu := (\lambda_{m+1} > \lambda_{m+2} > \dots > \lambda_r).$$

Then it holds that

$$\begin{aligned} GP_\lambda(x_1, x_2, \dots, x_n) &= \frac{1}{(n-m)!} \sum_{w \in S_n} w (A_\mu GP_\nu(x_{m+1}, x_{m+2}, \dots, x_n)), \\ GQ_\lambda(x_1, x_2, \dots, x_n) &= \frac{1}{(n-m)!} \sum_{w \in S_n} w (B_\mu GQ_\nu(x_{m+1}, x_{m+2}, \dots, x_n)). \end{aligned}$$

Proof. Choose any polynomial $f(x) \in \mathbb{Z}[\beta][x]$ and let $Z_i := f(x_i) \prod_{j=i+1}^n \frac{x_i \oplus x_j}{x_i \ominus x_j}$ for each $i \in [r]$. By Theorem 2.1, the expression

$$(3.2) \quad \frac{1}{(n-r)!} \sum_{w \in S_n} w (x^\lambda Z_1 Z_2 \cdots Z_r)$$

gives $GP_\lambda(x_1, \dots, x_n)$ when $f(x) = 1$ and $GQ_\lambda(x_1, \dots, x_n)$ when $f(x) = 2 + \beta x$. Let S_m and $H_{n-m} \cong S_{n-m}$ be the subgroups of permutations in S_n fixing each $i \in [n] \setminus [m]$ and $i \in [m]$ respectively. Then we can rewrite (3.2) as

$$\frac{1}{(n-r)!m!(n-m)!} \sum_{w \in S_n} w \left(\sum_{(g,h) \in S_m \times H_{n-m}} gh(x^\mu Z_1 \cdots Z_m) gh(x^{\tilde{\nu}} Z_{m+1} \cdots Z_r) \right)$$

where $\tilde{\nu}$ is the sequence formed from ν by prepending m zeros. The subgroups S_m and H_{n-m} commute, and each $h \in H_{n-m}$ fixes $x^\mu Z_1 \cdots Z_m$ while each $g \in S_m$ fixes $x^{\tilde{\nu}} Z_{m+1} \cdots Z_r$. The preceding expression is therefore equal to

$$\frac{1}{(n-r)!} \sum_{w \in S_n} w \left(\frac{1}{m!} \sum_{g \in S_m} g(x^\mu Z_1 \cdots Z_m) \cdot \frac{1}{(n-m)!} \sum_{h \in H_{n-m}} h(x^{\tilde{\nu}} Z_{m+1} \cdots Z_r) \right).$$

If $f(x) = 1$ then the internal sums here are $\frac{1}{m!} \sum_{g \in S_m} g(x^\mu Z_1 \cdots Z_m) = A_\mu$ and $\frac{1}{(n-m)!} \sum_{h \in H_{n-m}} h(x^{\tilde{\nu}} Z_{m+1} \cdots Z_r) = GP_\nu(x_{m+1}, \dots, x_n)$ by Theorem 2.1. This proves the first identity. The other follows by taking $f(x) = 2 + \beta x$. \square

For $r \in [n]$ let $\Pi^{r,n}(x) := \prod_{i=1}^r \prod_{j=i+1}^n \frac{x_i \oplus x_j}{x_i \ominus x_j}$.

LEMMA 3.2. Choose integers $1 \leq p < q \leq r$ and let $t_{pq} := (p, q) \in S_r$. Then

$$t_{pq} \left((1 + \beta x_p)^{q-p} \Pi^{r,n}(x) \right) = -(1 + \beta x_p)^{q-p} \Pi^{r,n}(x).$$

Consequently if $f(x) \in \mathbb{Z}[\beta][x_1, x_2, \dots]$ is any polynomial fixed by t_{pq} then

$$\sum_{w \in S_r} w \left(f(x) (1 + \beta x_p)^{q-p} \Pi^{r,n}(x) \right) = 0.$$

Proof. The first identity is a straightforward exercise in algebra. The second claim follows since $\sum_{w \in S_r} w = \frac{1}{2} \sum_{w \in S_r} w(1 + t_{pq}) \in \mathbb{Z}S_r$. \square

Choose an integer $m \in [n]$ and let $\delta := (m, m - 1, \dots, 3, 2, 1)$. For each $i \in [m]$ let $\mathbf{e}_i := (0, \dots, 0, 1, 0, \dots, 0)$ be the standard basis vector in \mathbb{Z}^m .

LEMMA 3.3. Let $v \in \mathbb{N}^m$. Then $A_{\delta+v} + \beta A_{\delta+v+\mathbf{e}_i} = 0$ whenever $v_{i+1} = v_i + 1$ for some $i \in [m - 1]$ or $v_{i+2} = v_{i+1} + 1 = v_i + 1$ for some $i \in [m - 2]$.


Proof. First assume $v_{i+1} = v_i + 1$ for some $i \in [m - 1]$. Then $x^{\delta+v}$ is a polynomial fixed by $t_{i,i+1}$ so $A_{\delta+v} + \beta A_{\delta+v+\mathbf{e}_i} = \frac{1}{m!} \sum_{w \in S_m} w(x^{\delta+v}(1 + \beta x_i) \Pi^{m,n}(x)) = 0$ by Lemma 3.2.

Next suppose that $v_{i+2} = v_{i+1} + 1 = v_i + 1$ for some $i \in [m - 2]$. Let $\alpha := \delta + v - \mathbf{e}_i$. Then we can write $\beta(A_{\delta+v} + \beta A_{\delta+v+\mathbf{e}_i})$ as

$$\frac{1}{m!} \sum_{w \in S_m} w(x^\alpha(1 + \beta x_i)^2 \Pi^{m,n}(x)) - \frac{1}{m!} \sum_{w \in S_m} w(x^\alpha(1 + \beta x_i) \Pi^{m,n}(x)).$$

Since x^α is fixed by $t_{i,i+2}$ and $t_{i,i+1}$, applying Lemma 3.2 with $(p, q) = (i, i + 2)$ and $(p, q) = (i, i + 1)$ shows that both terms are zero. The ring of rational functions in $\beta, x_1, x_2, \dots, x_n$ is an integral domain, so $A_{\delta+v} + \beta A_{\delta+v+\mathbf{e}_i} = 0$. \square

After expanding B_δ in terms of the A_λ 's, one can apply many cancellations from Lemma 3.3. We will use the next lemma to organize these cancellations. This lemma involves a certain directed graph \mathcal{G}_m for $m \geq 2$ which we define inductively. In general, the vertex set of \mathcal{G}_m consists of all nonempty subsets of $[m]$ excluding sets of the form $[i]$ for i odd and including two copies of $[i]$ for i even. When $m \in \{2, 3\}$ the graph \mathcal{G}_m is given explicitly by



Assume that $m \geq 4$ and that \mathcal{G}_{m-2} and \mathcal{G}_{m-1} have been constructed. Let \mathcal{A} be the set of vertices $S \in \mathcal{G}_m$ with $\{m - 2, m - 1, m\} \subseteq S$, let \mathcal{B} be the set of vertices $S \in \mathcal{G}_m$ with $m \notin S$, and let \mathcal{C} be the set of remaining vertices in \mathcal{G}_m . All of the doubled vertices $[i] \in \mathcal{G}_m$ for $i = 2, 4, 6, \dots$ belong to \mathcal{A} or \mathcal{B} .

The edges of \mathcal{G}_m are given as follows. The three sets \mathcal{A} , \mathcal{B} , and \mathcal{C} are each unions of connected components. An edge goes from $S \in \mathcal{A}$ to $T \in \mathcal{A}$ if and only if the edge $S \setminus \{m - 1, m\} \rightarrow T \setminus \{m - 1, m\}$ exists in \mathcal{G}_{m-2} . An edge goes from $S \in \mathcal{B}$ to $T \in \mathcal{B}$ if and only if the same edge $S \rightarrow T$ exists in \mathcal{G}_{m-1} . The elements of \mathcal{C} consist of the distinct unions $S \sqcup \{m - 1, m\}$, $S \sqcup \{m - 2, m\}$, and $S \sqcup \{m\}$ as S ranges over all subsets of $[m - 3]$, and for each $S \subseteq [m - 3]$ there are edges $S \sqcup \{m - 1, m\} \rightarrow S \sqcup \{m\}$ and $S \sqcup \{m - 2, m\} \rightarrow S \sqcup \{m\}$.

EXAMPLE 3.4. If $m = 4$ then

- \mathcal{A} has elements $\{1, 2, 3, 4\}, \{1, 2, 3, 4\}, \{2, 3, 4\};$
- \mathcal{B} has elements $\{1, 2\}, \{1, 2\}, \{3\}, \{1, 3\}, \{2, 3\}, \{2\};$

$(x_1 x_2 \dots x_{q-p+1})^{p-1}$ gives $B_\mu = 2^{q-p+1} A_\mu - 2^{q-p+1} \sum_{i=1}^{q-p+1} (-\beta/2)^i A_{\mu+e_{[i]}}$ which can be rewritten as (3.3). \square

If $\lambda = (\lambda_1 \geq \dots \geq \lambda_p > 0)$ and $\mu = (\mu_1 \geq \dots \geq \mu_q > 0)$ are partitions then let $\lambda\mu$ denote their concatenation; this will be another partition if $\lambda_p \geq \mu_1$. Given a strict partition μ , define $\Lambda(\mu)$ to be the set of strict partitions $\lambda \supseteq \mu$ with $\ell(\lambda) = \ell(\mu)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip. We can now prove Theorem 1.1, which states that $GQ_\mu = 2^{\ell(\mu)} \sum_{\lambda \in \Lambda(\mu)} (-1)^{\text{cols}(\lambda/\mu)} (-\beta/2)^{|\lambda/\mu|} GP_\lambda$.

Proof of Theorem 1.1. Let $\mu = (\mu_1 > \dots > \mu_r > 0)$ be a nonempty strict partition. We first prove the desired identity specialized to the variables x_1, x_2, \dots, x_n , so assume our fixed value of n has $n \geq r > 0$. Let $q = \mu_1$ and suppose $m \in [r]$ is maximal with $\mu_m = q + 1 - m$. We proceed by induction on $r - m$.

In the base case when $m = r$, the result to prove is Lemma 3.6. It remains to deal with the inductive step. Assume $1 \leq m < r$ and set

$$\gamma := (\mu_1 > \mu_2 > \dots > \mu_m) \quad \text{and} \quad \nu := (\mu_{m+1} > \mu_{m+2} > \dots > \mu_r).$$

Then $\gamma = (q, q - 1, q - 2, \dots, p)$ for $p := \mu_m$ and $\gamma_m \geq \nu_1 + 2$. We may assume by induction that the desired identity holds when μ is replaced by ν , since this replacement transforms $r \mapsto r - m$ and $m \mapsto$ (some positive number) so reduces the difference $r - m$. This assumption and Lemma 3.6 imply that $B_\gamma GQ_\nu(x_{m+1}, \dots, x_n)$ is equal to

$$2^r \sum_{\tilde{\gamma} \in \Lambda(\gamma)} \sum_{\tilde{\nu} \in \Lambda(\nu)} (-1)^{\text{cols}(\tilde{\gamma}/\gamma) + \text{cols}(\tilde{\nu}/\nu)} (-\beta/2)^{|\tilde{\gamma}/\gamma| + |\tilde{\nu}/\nu|} A_{\tilde{\gamma}} GP_{\tilde{\nu}}(x_{m+1}, \dots, x_n).$$

Using both parts of Lemma 3.1, we deduce that $GQ_\mu(x_1, \dots, x_n)$ is equal to

$$2^r \sum_{\tilde{\gamma} \in \Lambda(\gamma)} \sum_{\tilde{\nu} \in \Lambda(\nu)} (-1)^{\text{cols}(\tilde{\gamma}/\gamma) + \text{cols}(\tilde{\nu}/\nu)} (-\beta/2)^{|\tilde{\gamma}/\gamma| + |\tilde{\nu}/\nu|} GP_{\tilde{\gamma}\tilde{\nu}}(x_1, \dots, x_n).$$

Since $\mu = \gamma\nu$ and $\gamma_m \geq \nu_1 + 2$, the concatenation map $(\tilde{\gamma}, \tilde{\nu}) \mapsto \tilde{\gamma}\tilde{\nu}$ is a bijection $\Lambda(\gamma) \times \Lambda(\nu) \xrightarrow{\sim} \Lambda(\mu)$ and if $\lambda = \tilde{\gamma}\tilde{\nu}$ for $(\tilde{\gamma}, \tilde{\nu}) \in \Lambda(\gamma) \times \Lambda(\nu)$ then $\text{cols}(\tilde{\gamma}/\gamma) + \text{cols}(\tilde{\nu}/\nu) = \text{cols}(\lambda/\mu)$ and $|\tilde{\gamma}/\gamma| + |\tilde{\nu}/\nu| = |\lambda/\mu|$. Hence

$$GQ_\mu(x_1, \dots, x_n) = 2^r \sum_{\lambda \in \Lambda(\mu)} (-1)^{\text{cols}(\lambda/\mu)} (-\beta/2)^{|\lambda/\mu|} GP_\lambda(x_1, \dots, x_n).$$

This even holds when $\mu = \emptyset$, so taking the limit as $n \rightarrow \infty$ gives the theorem. \square

4. WEIGHT-PRESERVING BIJECTIONS

As the GQ_λ 's and GP_λ 's are generating functions for set-valued shifted tableaux, Theorem 1.1 has some enumerative consequences, which we describe here.

Let $\mathcal{X} \subseteq \{1, 2, 3, \dots\} \times \{1, 2, 3, \dots\}$ be a set of positions. Given a set-valued shifted tableau T , define $\text{unprime}_{\max}^{\mathcal{X}}(T)$ to be the tableau formed from T by removing the prime from the largest element of T_{ij} for each $(i, j) \in \mathcal{X}$, whenever this element is not already primed. If $\mathcal{X} = \{(1, 1), (1, 2), (1, 3)\}$ then

$$\text{unprime}_{\max}^{\mathcal{X}} \left(\begin{array}{|c|c|c|} \hline & 34' & \\ \hline 2'2 & 3' & 34'6' \\ \hline \end{array} \right) = \begin{array}{|c|c|c|} \hline & 34' & \\ \hline 2'2 & 3 & 34'6 \\ \hline \end{array}$$

for example. If T is semistandard and $\mathcal{X} \subseteq \{(1, 1), (2, 2), (3, 3), \dots\}$ then $\text{unprime}_{\max}^{\mathcal{X}}(T)$ is also semistandard. This property may fail if \mathcal{X} is not a subset of the main diagonal (as we see in the previous example).

If λ is a strict partition and $\mathcal{X} \subseteq \{(i, i) : i \in [\ell(\lambda)]\}$ is a set of diagonal positions, then each $T \in \text{SetShYT}_P(\lambda)$ has exactly $2^{|\mathcal{X}|}$ preimages in $\text{SetShYT}_Q(\lambda)$ under $\text{unprime}_{\max}^{\mathcal{X}}$. Given strict partitions $\lambda \supseteq \mu$ define $\text{SetShYT}_P(\lambda : \mu)$ to be the set of semistandard set-valued shifted tableaux $T \in \text{SetShYT}_Q(\lambda)$ with

$$\text{unprime}_{\max}^{\mathcal{X}}(T) \in \text{SetShYT}_P(\lambda) \text{ for } \mathcal{X} := \{(i, i) : \lambda_i = \mu_i\}.$$

The diagonal entry in row i of such a tableau can have at most one primed element if $\lambda_i = \mu_i$ and no primed elements if $\lambda_i > \mu_i$.

Finally, for a strict partition μ let $\Lambda^{\pm}(\mu)$ be the set of strict partitions $\lambda \supseteq \mu$ with $\ell(\lambda) = \ell(\mu)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip and $(-1)^{\text{cols}(\lambda/\mu)+|\lambda/\mu|} = \pm 1$. Then $\Lambda(\mu) = \Lambda^+(\mu) \sqcup \Lambda^-(\mu)$ and this decomposition reflects the decomposition of the right side of Theorem 1.1 into positive and negative terms.

COROLLARY 4.1. *For each strict partition μ there is a weight-preserving bijection*

$$\text{SetShYT}_Q(\mu) \sqcup \bigsqcup_{\lambda \in \Lambda^-(\mu)} \text{SetShYT}_P(\lambda : \mu) \rightarrow \bigsqcup_{\lambda \in \Lambda^+(\mu)} \text{SetShYT}_P(\lambda : \mu).$$

Proof. To see that the domain of the given map is indeed a disjoint union, observe that the set $\Lambda^-(\mu)$ is empty if and only if all parts of μ differ by at least two. Since $\text{cols}(\mu/\mu) + |\mu/\mu| = 0$, the set $\Lambda^+(\mu)$ is never empty and $\text{SetShYT}_Q(\mu)$ and $\text{SetShYT}_P(\lambda : \mu)$ are disjoint for all $\lambda \in \Lambda^-(\mu)$.

If $\lambda \in \Lambda^{\pm}(\mu)$ then $\ell(\mu) - |\lambda/\mu| = |\{i : \lambda_i = \mu_i\}|$. Thus, moving the negative terms on the right side of (1.4) to the left side, then substituting the generating functions for GQ_{μ} and GP_{λ} in (1.3), and finally setting $\beta = 1$ gives

$$(4.1) \quad \sum_{T \in \text{SetShYT}_Q(\mu)} x^T + \sum_{\lambda \in \Lambda^-(\mu)} \sum_{T \in \text{SetShYT}_P(\lambda)} 2^{|\{i : \lambda_i = \mu_i\}|} x^T$$

is equal to

$$(4.2) \quad \sum_{\lambda \in \Lambda^+(\mu)} \sum_{T \in \text{SetShYT}_P(\lambda)} 2^{|\{i : \lambda_i = \mu_i\}|} x^T.$$

We have $\sum_{T \in \text{SetShYT}_P(\lambda)} 2^{|\{i : \lambda_i = \mu_i\}|} x^T = \sum_{T \in \text{SetShYT}_P(\lambda : \mu)} x^T$ for any strict partition λ , and the corollary follows by substituting this identity into (4.1) and (4.2) and equating coefficients. \square

It is an interesting open problem to find a bijective proof of Theorem 1.1. One way to achieve this would be to construct an explicit map realizing Corollary 4.1. This is easy to do when $\mu = (n)$ has only one nonzero part, in which case the bijection in Corollary 4.1 is a map

$$(4.3) \quad \text{SetShYT}_Q(n) \rightarrow \text{SetShYT}_P(n : n) \sqcup \text{SetShYT}_P(n + 1).$$

The set $\text{SetShYT}_P(n : n)$ is contained in $\text{SetShYT}_Q(n)$ and is the union of $\text{SetShYT}_P(n)$ and the set of tableaux formed from elements of $\text{SetShYT}_P(n)$ by adding a prime to the largest number in box $(1, 1)$.

A bijection (4.3) is given by mapping each $T \in \text{SetShYT}_P(n : n)$ to itself and each $T \in \text{SetShYT}_Q(n) \setminus \text{SetShYT}_P(n : n)$ to the tableau in $\text{SetShYT}_P(n + 1)$ formed by adding $\frac{1}{2}$ to the smallest primed number $i' = i - \frac{1}{2}$ in box $(1, 1)$, and then splitting this diagonal box into two adjacent boxes containing all numbers $\leq i$ and $> i$, respectively. This map is weight-preserving and would send

$$\begin{array}{|c|c|c|} \hline 12'3'3 & 34 & 5' \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|} \hline 12 & 3'3 & 34 & 5' \\ \hline \end{array}$$

for example. It seems difficult to generalize this idea to larger shapes. Even in the next simplest case $\mu = (3, 1)$ we do not know of a straightforward way to describe a weight-preserving bijection of the form in Corollary 4.1.

In [8, §9.2], Ikeda and Naruse derive another set of combinatorial formulas for GP_λ and GQ_λ as generating functions for *excited Young diagrams*. One could also try to find a bijective proof of Theorem 1.1 via these expressions.

5. SKEW ANALOGUES

Let $\mu \subseteq \lambda$ be strict partitions. A *semistandard (skew) shifted tableau* of shape λ/μ is a filling of $SD_{\lambda/\mu} := SD_\lambda \setminus SD_\mu$ by positive half-integers such that rows and columns are weakly increasing, with no primed entries repeated in a row and no unprimed entries repeated in a column.

Let $ShYT_Q(\lambda/\mu)$ be the set of all such tableaux and let $ShYT_P(\lambda/\mu)$ be the subset in which primed entries are disallowed from diagonal positions. We define both sets to be empty if $\mu \not\subseteq \lambda$. The *skew Schur P- and Q-functions* are

$$(5.1) \quad P_{\lambda/\mu} := \sum_{T \in ShYT_P(\lambda/\mu)} x^T \quad \text{and} \quad Q_{\lambda/\mu} := \sum_{T \in ShYT_Q(\lambda/\mu)} x^T$$

where as usual $x^T := \prod_{i \geq 1} x_i^{m_i}$ with m_i denoting the number of entries of T equal to i or i' . To motivate these symmetric functions, we need some additional notation. If $f \in \mathbb{Z}[\beta][[x_1, x_2, \dots]]$, then write $f(x, y)$ for the power series $f(x_1, y_1, x_2, y_2, \dots)$ where x_1, x_2, \dots and y_1, y_2, \dots are separate sets of commuting variables; we also set $f(x) := f(x_1, x_2, \dots) = f$ and $f(y) := f(y_1, y_2, \dots)$. If f is symmetric then specializing $f(x, y)$ to finitely many variables gives

$$f(x_1, y_1, x_2, y_2, \dots, x_n, y_n) = f(x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n).$$

It follows that we can write

$$(5.2) \quad P_\lambda(x, y) = \sum_\nu P_\nu(x)P_{\lambda/\nu}(y) \quad \text{and} \quad Q_\lambda(x, y) = \sum_\nu Q_\nu(x)Q_{\lambda/\nu}(y)$$

where in both sums ν ranges over all strict partitions [22, Eq. (8.2)].

Define the set $SetShYT_Q(\lambda/\mu)$ of *semistandard set-valued (skew) shifted tableaux* of shape λ/μ in the same way as $SetShYT_Q(\lambda)$, just replacing references to “fillings of SD_λ ” by “fillings of $SD_{\lambda/\mu}$.” Let $SetShYT_P(\lambda/\mu)$ be the subset of tableaux in $SetShYT_Q(\lambda/\mu)$ with no primed numbers in any diagonal boxes. The *skew K-theoretic Schur P- and Q-functions* are then

$$(5.3) \quad \begin{aligned} GP_{\lambda/\mu} &:= \sum_{T \in SetShYT_P(\lambda/\mu)} \beta^{|T| - |\lambda/\mu|} x^T, \\ GQ_{\lambda/\mu} &:= \sum_{T \in SetShYT_Q(\lambda/\mu)} \beta^{|T| - |\lambda/\mu|} x^T, \end{aligned}$$

where x^T is defined in the same way as for elements of $SetShYT_Q(\lambda)$. When $\mu \not\subseteq \lambda$ we consider both $SetShYT_P(\lambda/\mu)$ and $SetShYT_Q(\lambda/\mu)$ to be empty so that $GP_{\lambda/\mu} = GQ_{\lambda/\mu} = 0$. These generalizations of GP_λ and GQ_λ were first defined in [11, §4.6] in the context of *enriched set-valued P-partitions*.

The K -theoretic version of (5.2) involves a variant of these power series. The *removable boxes* of μ are the positions $(i, j) \in SD_\mu$ such that $SD_\mu \setminus \{(i, j)\}$ is the shifted diagram of another strict partition. Let $Rem(\mu)$ be the set of removable boxes of the

strict partition μ . For strict partitions $\mu \subseteq \lambda$ define

$$(5.4) \quad \begin{aligned} GP_{\lambda//\mu} &:= \sum_{\nu \subseteq \mu, \text{SD}_{\mu/\nu} \subseteq \text{Rem}(\mu)} \beta^{|\mu/\nu|} GP_{\lambda/\nu}, \\ GQ_{\lambda//\mu} &:= \sum_{\nu \subseteq \mu, \text{SD}_{\mu/\nu} \subseteq \text{Rem}(\mu)} \beta^{|\mu/\nu|} GQ_{\lambda/\nu}, \end{aligned}$$

where in both sums ν must be a strict partition. For strict partitions $\mu \not\subseteq \lambda$ we set $GP_{\lambda//\mu} = GQ_{\lambda//\mu} = 0$. Then the definitions of GP_λ and GQ_λ as set-valued shifted tableau generating functions imply that

$$(5.5) \quad \begin{aligned} GP_\lambda(x, y) &= \sum_\nu GP_\nu(x) GP_{\lambda//\nu}(y), \\ GQ_\lambda(x, y) &= \sum_\nu GQ_\nu(x) GQ_{\lambda//\nu}(y), \end{aligned}$$

where both sums are over all strict partitions ν .

Since the GP_ν 's and GQ_ν 's are linearly independent and symmetric, these identities imply that $GP_{\lambda//\mu}$ and $GQ_{\lambda//\mu}$ are symmetric. Hence $GP_{\lambda/\mu}$ and $GQ_{\lambda/\mu}$ are also symmetric as they can be written in terms of $GP_{\lambda//\mu}$ and $GQ_{\lambda//\mu}$ via inclusion-exclusion [11, Cor. 5.7].

More strongly, $GQ_{\lambda//\mu}$ (respectively, $GP_{\lambda//\mu}$) is a possibly infinite $\mathbb{Z}[\beta]$ -linear combination of GQ -functions (respectively, GP -functions) by [11, Cor 5.13]. Since both functions, when nonzero, are homogeneous of degree $|\lambda| - |\mu|$ if we set $\deg(\beta) = -1$ and $\deg(x_i) = 1$, it follows that

$$(5.6) \quad \begin{aligned} GQ_{\lambda//\mu} &= \sum_\nu \hat{a}_{\mu\nu}^\lambda \beta^{|\mu|+|\nu|-|\lambda|} GQ_\nu, \\ GP_{\lambda//\mu} &= \sum_\nu \hat{b}_{\mu\nu}^\lambda \beta^{|\mu|+|\nu|-|\lambda|} GP_\nu, \end{aligned}$$

for unique integers $\hat{a}_{\mu\nu}^\lambda, \hat{b}_{\mu\nu}^\lambda \in \mathbb{Z}$ which must be zero when $|\mu| + |\nu| < |\lambda|$. It is expected that these coefficients are all nonnegative, and nonzero for only finitely many strict partitions ν ; see [11, Conj. 5.14].

Given strict partitions $\mu \subseteq \lambda$, we consider the statistic

$$\text{overlap}(\lambda/\mu) := |\{(i, j) \in \text{SD}_{\lambda/\mu} : (i-1, j) \in \text{SD}_{\lambda/\mu}\}|.$$

This quantity is closely related to $\text{cols}(\lambda/\mu) := |\{j : (i, j) \in \text{SD}_{\lambda/\mu}\}|$.

LEMMA 5.1. *Let $\mu \subseteq \lambda$ be strict partitions with $\ell(\mu) = \ell(\lambda)$. Then*

$$\sum_\eta (-1)^{\text{cols}(\lambda/\eta)} 2^{\text{overlap}(\eta/\mu)} = \sum_\gamma (-1)^{\text{cols}(\gamma/\mu)} 2^{\text{overlap}(\lambda/\gamma)} = \begin{cases} 1 & \text{if } \mu = \lambda \\ 0 & \text{if } \mu \neq \lambda \end{cases}$$

where the first summation is over all strict partitions η with $\mu \subseteq \eta \subseteq \lambda$ such that $\text{SD}_{\lambda/\eta}$ is a vertical strip, and the second summation is over all strict partitions γ with $\mu \subseteq \gamma \subseteq \lambda$ such that $\text{SD}_{\gamma/\mu}$ is a vertical strip.

Proof. The desired identity is clear if $\mu = \lambda$. Assume $\mu \neq \lambda$ so that $\text{SD}_{\lambda/\mu}$ is nonempty. We start by showing that the first sum is zero. Suppose the rightmost box of $\text{SD}_{\lambda/\mu}$ is in column n and $\text{SD}_{\lambda/\mu}$ contains $k > 0$ boxes in this column. Choose a strict partition η with $\mu \subseteq \eta \subseteq \lambda$ such that $\text{SD}_{\lambda/\eta}$ is a vertical strip. Let $L := \{(i, j) \in \text{SD}_{\eta/\mu} : j < n\}$ and $R := \{(i, j) \in \text{SD}_{\eta/\mu} : j = n\}$ so that $\text{SD}_{\eta/\mu} = L \sqcup R$. Because $\text{SD}_{\lambda/\eta}$ is a vertical strip, there are only $k+1$ possibilities for R , which must be a set of adjacent positions

at the bottom of column n in $\text{SD}_{\lambda/\mu}$. Moreover, when L is fixed and η varies, each of these possibilities for R occurs exactly once. Now observe that if $r := |R|$ then

$$\begin{aligned} \text{overlap}(\eta/\mu) &= |\{(i, j) \in L : (i - 1, j) \in L\}| + |\{(i, j) \in R : (i - 1, j) \in R\}| \\ &= |\{(i, j) \in L : (i - 1, j) \in L\}| + \max\{0, r - 1\} \end{aligned}$$

and also

$$\text{cols}(\lambda/\eta) = |\{j : (i, j) \in \text{SD}_{\lambda/\eta} \sqcup R\}| - \begin{cases} 1 & \text{if } r = k \\ 0 & \text{if } 0 \leq r < k. \end{cases}$$

Since $\text{SD}_{\lambda/\eta} \sqcup R = \text{SD}_{\lambda/\mu} - L$, we can rewrite the preceding identity as

$$\text{cols}(\lambda/\eta) = |\{j : (i, j) \in \text{SD}_{\lambda/\mu} - L\}| + \min\{0, k - r - 1\}.$$

By substituting these formulas and factoring out the terms depending on L , we deduce that the sum $\sum_{\eta} (-1)^{\text{cols}(\lambda/\eta)} 2^{\text{overlap}(\eta/\mu)}$ is a multiple of

$$(5.7) \quad \sum_{r=0}^k (-1)^{\min\{0, k-r-1\}} 2^{\max\{0, r-1\}} = 2^0 + 2^0 + 2^1 + \dots + 2^{k-2} - 2^{k-1} = 0$$

and so is zero itself.

A similar argument shows that the other sum in the lemma is zero. Suppose now that the *leftmost* box of $\text{SD}_{\lambda/\mu}$ is in column n and $\text{SD}_{\lambda/\mu}$ contains $k > 0$ boxes in this column. Choose a strict partition γ with $\mu \subseteq \gamma \subseteq \lambda$ such that $\text{SD}_{\gamma/\mu}$ is a vertical strip. Let $L := \{(i, j) \in \text{SD}_{\lambda/\gamma} : j = n\}$ and $R := \{(i, j) \in \text{SD}_{\lambda/\gamma} : j > n\}$ so that $\text{SD}_{\lambda/\gamma} = L \sqcup R$. Because $\text{SD}_{\gamma/\mu}$ is a vertical strip, there are now only $k + 1$ possibilities for L , which must be a set of adjacent positions at the top of column n in $\text{SD}_{\lambda/\mu}$, and when R is fixed and γ varies, each of these possibilities occurs exactly once. Finally, if $\ell := |L|$ then we have

$$\begin{aligned} \text{overlap}(\lambda/\gamma) &= |\{(i, j) \in R : (i - 1, j) \in R\}| + |\{(i, j) \in L : (i - 1, j) \in L\}| \\ &= |\{(i, j) \in R : (i - 1, j) \in R\}| + \max\{0, \ell - 1\} \end{aligned}$$

and also

$$\text{cols}(\gamma/\mu) = |\{j : (i, j) \in \text{SD}_{\gamma/\mu} \sqcup L\}| - \begin{cases} 1 & \text{if } \ell = k \\ 0 & \text{if } 0 \leq \ell < k. \end{cases}$$

Since $\text{SD}_{\gamma/\mu} \sqcup L = \text{SD}_{\lambda/\mu} - R$, we can rewrite the preceding identity as

$$\text{cols}(\gamma/\mu) = |\{j : (i, j) \in \text{SD}_{\lambda/\mu} - R\}| + \min\{0, k - \ell - 1\}.$$

By substituting these formulas and factoring out the terms depending on R , we deduce that $\sum_{\gamma} (-1)^{\text{cols}(\gamma/\mu)} 2^{\text{overlap}(\lambda/\gamma)}$ is also a multiple of (5.7), as needed. \square

REMARK 5.2. For strict partitions λ and μ , define $M_{\lambda\mu} := 0$ when $\mu \not\subseteq \lambda$ or $\ell(\mu) \neq \ell(\lambda)$, and otherwise set $M_{\lambda\mu} := 2^{\text{overlap}(\lambda/\mu)}$. Also define $N_{\lambda\mu} := 0$ when $\mu \not\subseteq \lambda$ or $\ell(\mu) \neq \ell(\lambda)$ or $\text{SD}_{\lambda/\mu}$ is not a vertical strip, and otherwise set $N_{\lambda\mu} := (-1)^{\text{cols}(\lambda/\mu)}$. Lemma 5.1 asserts that the matrices $[M_{\lambda\mu}]$ and $[N_{\lambda\mu}]$ are inverses.

We can now derive a skew generalization of Theorem 1.1.

THEOREM 5.3. *Suppose $\nu \subseteq \mu$ are strict partitions. Then*

$$GQ_{\mu//\nu} = \sum_{(\kappa, \lambda)} 2^{\ell(\mu) - \ell(\nu) + \text{overlap}(\nu/\kappa)} (-1)^{\text{cols}(\lambda/\mu)} (-\beta/2)^{|\nu/\kappa| + |\lambda/\mu|} GP_{\lambda//\kappa}$$

where the sum is over all pairs of strict partitions (κ, λ) with $\kappa \subseteq \nu \subseteq \mu \subseteq \lambda$ and $\ell(\kappa) = \ell(\nu) \leq \ell(\mu) = \ell(\lambda)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip.

Proof. Expanding $GQ_\mu(x, y)$ using Theorem 1.1 and then applying (5.5) gives

$$GQ_\mu(x, y) = \sum_\lambda \sum_\nu 2^{\ell(\mu)} (-1)^{\text{cols}(\lambda/\mu)} (-\beta/2)^{|\lambda/\mu|} GP_\nu(x) GP_{\lambda//\nu}(y)$$

where the first sum is over strict partitions $\lambda \supseteq \mu$ with $\ell(\lambda) = \ell(\mu)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip and the second sum is over all strict partitions $\nu \subseteq \lambda$. Alternatively, using Theorem 1.1 to expand the right side of (5.5) gives

$$GQ_\mu(x, y) = \sum_\eta \sum_\nu 2^{\ell(\nu)} (-1)^{\text{cols}(\nu/\eta)} (-\beta/2)^{|\nu/\eta|} GP_\nu(x) GQ_{\mu//\eta}(y)$$

where the first sum is over all strict partitions $\eta \subseteq \mu$ and the second sum over strict partitions $\nu \supseteq \eta$ with $\ell(\nu) = \ell(\eta)$ such that $\text{SD}_{\nu/\eta}$ is a vertical strip.⁽¹⁾ Equating the coefficients of GP_ν in these expressions gives

$$\sum_\eta 2^{\ell(\nu)} (-1)^{\text{cols}(\nu/\eta)} \left(\frac{-\beta}{2}\right)^{|\nu/\eta|} GQ_{\mu//\eta} = \sum_\lambda 2^{\ell(\mu)} (-1)^{\text{cols}(\lambda/\mu)} \left(\frac{-\beta}{2}\right)^{|\lambda/\mu|} GP_{\lambda//\nu}$$

where the sums are over certain strict partitions η and λ ; to be precise, assuming $\nu \subseteq \mu$, the preceding equation is equivalent to

$$GQ_{\mu//\nu} = \sum_\lambda 2^{\ell(\mu)-\ell(\nu)} (-1)^{\text{cols}(\lambda/\mu)} \left(\frac{-\beta}{2}\right)^{|\lambda/\mu|} GP_{\lambda//\nu} - \sum_{\eta \subsetneq \nu} (-1)^{\text{cols}(\nu/\eta)} \left(\frac{-\beta}{2}\right)^{|\nu/\eta|} GQ_{\mu//\eta}$$

where the first sum is over strict partitions $\lambda \supseteq \mu$ with $\ell(\lambda) = \ell(\mu)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip and the second sum is over strict partitions $\eta \subsetneq \nu$ with $\ell(\eta) = \ell(\nu)$ such that $\text{SD}_{\nu/\eta}$ is a vertical strip.

When $\nu = (m, m-1, \dots, 2, 1)$ for some $m \in \mathbb{N}$, the sum over η has zero terms and the preceding formula reduces to the desired identity. Otherwise, we may assume by induction that the desired formula holds for each $GQ_{\mu//\eta}$. Substituting these formulas into the displayed equation gives an expression for $GQ_{\mu//\nu}$ as a linear combination of $GP_{\lambda//\kappa}$'s where κ and λ range over all strict partitions with $\kappa \subseteq \nu \subseteq \mu \subseteq \lambda$ and $\ell(\kappa) = \ell(\nu) \leq \ell(\mu) = \ell(\lambda)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip. The coefficient of $GP_{\lambda//\nu}$ in this expansion is $2^{\ell(\mu)-\ell(\nu)} (-1)^{\text{cols}(\lambda/\mu)} \left(\frac{-\beta}{2}\right)^{|\lambda/\mu|}$ as desired. The coefficient of $GP_{\lambda//\kappa}$ when $\kappa \subsetneq \nu \subseteq \mu \subseteq \lambda$ and $\ell(\kappa) = \ell(\nu) \leq \ell(\mu) = \ell(\lambda)$ and $\text{SD}_{\lambda/\mu}$ is a vertical strip is

$$- \sum_\eta (-1)^{\text{cols}(\nu/\eta)} \left(\frac{-\beta}{2}\right)^{|\nu/\eta|} \left(2^{\ell(\mu)-\ell(\eta)+\text{overlap}(\eta/\kappa)} (-1)^{\text{cols}(\lambda/\mu)} \left(\frac{-\beta}{2}\right)^{|\eta/\kappa|+|\lambda/\mu|} \right)$$

where the sum is over all strict partitions η with $\kappa \subseteq \eta \subsetneq \nu$ such that $\text{SD}_{\nu/\eta}$ is a vertical strip. Since then $\ell(\kappa) = \ell(\eta) = \ell(\nu)$, we can rewrite this as

$$2^{\ell(\mu)-\ell(\nu)} (-1)^{\text{cols}(\lambda/\mu)} \left(\frac{-\beta}{2}\right)^{|\lambda/\mu|+|\nu/\kappa|} \sum_\eta (-1)^{\text{cols}(\nu/\eta)+1} 2^{\text{overlap}(\eta/\kappa)},$$

so it suffices to show that $\sum_\eta (-1)^{\text{cols}(\nu/\eta)+1} 2^{\text{overlap}(\eta/\kappa)} = 2^{\text{overlap}(\nu/\kappa)}$ where the sum is again over η with $\kappa \subseteq \eta \subsetneq \nu$ such that $\text{SD}_{\nu/\eta}$ is a vertical strip. After moving all terms to one side, this identity is half of Lemma 5.1. \square

We mention one corollary, which was noted in passing above. Let $\delta_m := (m, m-1, \dots, 2, 1)$ for $m \in \mathbb{N}$. The following recovers Theorem 1.1 when $m = 0$.

⁽¹⁾When using Theorem 1.1 to expand $GQ_\eta(x)$ in $GQ_\mu(x, y) = \sum_\eta GQ_\eta(x) GQ_{\mu//\eta}(y)$ one expects to see the factor $2^{\ell(\eta)}$, but this can be changed to $2^{\ell(\nu)}$ since $\ell(\nu) = \ell(\eta)$.

COROLLARY 5.4. *Suppose μ is a strict partition with $\delta_m \subseteq \mu$. Then*

$$GQ_{\mu//\delta_m} = 2^{\ell(\mu)-m} \sum_{\lambda} (-1)^{\text{cols}(\lambda/\mu)} (-\beta/2)^{|\lambda/\mu|} GP_{\lambda//\delta_m}$$


where the sum is over strict partitions $\lambda \supseteq \mu$ with $\ell(\lambda) = \ell(\mu)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip.

Proof. If $\nu = \delta_m$ in Theorem 5.3, then the only strict partition $\kappa \subseteq \nu$ with $\ell(\kappa) = \ell(\nu)$ is $\kappa = \nu = \delta_m$, which has $\text{overlap}(\nu/\kappa) = |\nu/\kappa| = 0$. \square

There is some interest in determining when there are coincidences $P_{\lambda/\mu} = P_{\nu/\kappa}$ and $Q_{\lambda/\mu} = Q_{\nu/\kappa}$ among the skew Schur P - and Q -functions [1, 5, 21]. This phenomenon is less well-understood than for skew Schur functions.

One could consider the same problem for the skew GP - and GQ -functions. In particular, any equality $GP_{\lambda/\mu} = GP_{\nu/\kappa}$ would imply that $P_{\lambda/\mu} = P_{\nu/\kappa}$ (and likewise for the Q -functions), but it is not clear if the converse always holds. With these questions in mind, we explain one nontrivial equality between Schur Q -functions that generalizes to the K -theoretic setting.

Define the *flip* of a skew shape λ/μ to be the skew shape $\phi(\lambda/\mu)$ whose shifted diagram $\text{SD}_{\phi(\lambda/\mu)}$ is formed by reflecting $\text{SD}_{\lambda/\mu}$ across a line perpendicular to the main diagonal, so that in French notation the bottom row becomes the rightmost column. We refer to this operation as “flipping the diagram $\text{SD}_{\lambda/\mu}$ ”:



The following reduces to [4, Prop. IV.13] when $\beta = 0$:

PROPOSITION 5.5. *Let $\mu \subseteq \lambda$ be strict partitions, then*

$$GP_{\phi(\lambda/\mu)} = GP_{\lambda/\mu} \quad \text{and} \quad GQ_{\phi(\lambda/\mu)} = GQ_{\lambda/\mu}.$$

Proof. We first prove the GQ -identity. Suppose $T \in \text{SetShYT}_Q(\lambda/\mu)$. Write $\min(T)$ and $\max(T)$ for the minimal and the maximal numbers appearing in any entry of T . Let $n := \lceil \min(T) \rceil$ be whichever of $\min(T)$ or $\min(T) + \frac{1}{2}$ is an integer, and define $N := \lceil \max(T) \rceil$ similarly. Now let $\phi_Q(T)$ be the set-valued shifted tableau of shape $\phi(\lambda/\mu)$ formed by flipping T and then replacing each number a in each set-valued entry by $n + N - \frac{1}{2} - a$.

The rows and columns of $\phi_Q(T)$ are weakly increasing since the rows and columns of T are weakly increasing. As exactly one of a or $n + N - \frac{1}{2} - a$ is primed, the tableau $\phi_Q(T)$ does not have any unprimed numbers repeated in a column or primed numbers repeated in a row. We conclude that ϕ_Q defines a map $\text{SetShYT}_Q(\lambda/\mu) \rightarrow \text{SetShYT}_Q(\phi(\lambda/\mu))$, which is clearly invertible.

Notice that the weight of $\phi_Q(T)$ is $x^{\phi_Q(T)} = \sigma(x^T)$, where σ is the permutation with $\sigma(a) = n + N - a$ for all $n \leq a \leq N$. Since the values of n and N are determined by the monomial x^T , the symmetry of $GQ_{\lambda/\mu}$ implies that

$$(5.8) \quad GQ_{\lambda/\mu} = \sum_{T \in \text{SetShYT}_Q(\lambda/\mu)} \beta^{|\phi_Q(T)| - |\lambda/\mu|} x^{\phi_Q(T)} = GQ_{\phi(\lambda/\mu)}.$$

The proof of the GP -identity is similar, except now for $T \in \text{SetShYT}_P(\lambda/\mu)$ we define $\phi_P(T)$ from $\phi_Q(T)$ by adding $\frac{1}{2}$ to all numbers in diagonal positions. Since all numbers in diagonal entries of $\phi_Q(T)$ are primed when $T \in \text{SetShYT}_P(\lambda/\mu)$,

we have $x^{\phi_P(T)} = x^{\phi_Q(T)}$ and it follows that ϕ_P is a bijection $\text{SetShYT}_P(\lambda/\mu) \rightarrow \text{SetShYT}_P(\phi(\lambda/\mu))$. We can therefore replace every letter “ Q ” in (5.8) by “ P ” to deduce that $GP_{\lambda/\mu} = GP_{\phi(\lambda/\mu)}$. \square

It is not clear if there is a meaningful way to extend the preceding result to $GP_{\lambda//\mu}$ and $GQ_{\lambda//\mu}$. If one defines $\phi(\lambda//\mu) := \tilde{\lambda}//\tilde{\mu}$ where $\phi(\lambda/\mu) = \tilde{\lambda}/\tilde{\mu}$, for example, then it may hold that $GP_{\phi(\lambda//\mu)} \neq GP_{\lambda//\mu}$ and $GQ_{\phi(\lambda//\mu)} \neq GQ_{\lambda//\mu}$.

6. DUAL FUNCTIONS

Let $\bar{x} := \frac{-x}{1+\beta x}$ so that $x \oplus \bar{x} = 0$. Nakagawa and Naruse [16] define the *dual K -theoretic Schur P - and Q -functions* gp_λ and gq_λ to be the unique elements of $\mathbb{Z}[\beta][[x_1, x_2, \dots]]$ indexed by strict partitions λ such that

$$(6.1) \quad \sum_{\lambda} GQ_{\lambda}(x)gp_{\lambda}(y) = \sum_{\lambda} GP_{\lambda}(x)gq_{\lambda}(y) = \prod_{i,j \geq 1} \frac{1 - \bar{x}_i y_j}{1 - x_i y_j}.$$

Both sums in this *Cauchy identity* are over all strict partitions λ .

The power series gp_λ and gq_λ are a special case of the *dual universal factorial Schur P - and Q -functions* in [16, Def. 3.2]. One reason these specialization are interesting (compared to the more general “universal” functions) is because they have conjectural formulas as generating functions for *shifted reverse plane partitions* [16, Conj 5.1]. We will discuss this idea in Section 7.

Both gp_λ and gq_λ are symmetric in the x_i variables and of degree $|\lambda|$ if we set $\deg(\beta) = 0$ and $\deg(x_i) = 1$ [16, §3.2]. The sets $\{gp_\lambda : \lambda \text{ is a strict partition}\}$ and $\{gq_\lambda : \lambda \text{ is a strict partition}\}$ are $\mathbb{Z}[\beta]$ -bases for subrings of $\mathbb{Z}[\beta][[x_1, x_2, \dots]]$ by [16, Thm. 3.1], and it holds that $P_\lambda = gp_\lambda|_{\beta=0}$ and $Q_\lambda = gq_\lambda|_{\beta=0}$ [16, §3.2].

PROPOSITION 6.1. *We recover gp_λ from $gp_\lambda|_{\beta=1}$ (respectively, gq_λ from $gq_\lambda|_{\beta=1}$) by substituting $x_i \mapsto \beta^{-1}x_i$ for all i and then multiplying by $\beta^{|\lambda|}$. As such, if we set $\deg(\beta) = \deg(x_i) = 1$ then gp_λ and gq_λ are homogeneous of degree $|\lambda|$.*

Proof. We recover the original form of $\prod_{i,j \geq 1} \frac{1 - \bar{x}_i y_j}{1 - x_i y_j}$ after setting $\beta = 1$ by substituting $x_i \mapsto \beta x_i$ and $y_j \mapsto \beta^{-1}y_j$ for all i and j . It follows that if we define $gp_\lambda^{(1)} := gp_\lambda|_{\beta=1}$ and $GQ_\lambda^{(1)} := GQ_\lambda|_{\beta=1}$ then

$$\begin{aligned} \sum_{\lambda} GQ_{\lambda}^{(1)}(\beta x_1, \beta x_2, \dots)gp_{\lambda}^{(1)}(\beta^{-1}y_1, \beta^{-1}y_2, \dots) &= \prod_{i,j \geq 1} \frac{1 - \bar{x}_i y_j}{1 - x_i y_j} \\ &= \sum_{\lambda} GQ_{\lambda}(x)gp_{\lambda}(y). \end{aligned}$$

As it is clear from (1.3) that $GQ_\lambda(x) = \beta^{-|\lambda|}GQ_\lambda^{(1)}(\beta x_1, \beta x_2, \dots)$, this equation can only hold if $gp_\lambda(y) = \beta^{|\lambda|}gp_\lambda^{(1)}(\beta^{-1}y_1, \beta^{-1}y_2, \dots)$ which is equivalent to what is claimed in the lemma. The identity for gq_λ follows similarly. \square

Theorem 1.1 has a dual version that gives a *gp*-expansion of gq_λ :

COROLLARY 6.2. *If λ is a strict partition then*

$$gq_\lambda = 2^{\ell(\lambda)} \sum_{\mu} (-1)^{\text{cols}(\lambda/\mu)} (-\beta/2)^{|\lambda/\mu|} gp_\mu$$

where the sum is over strict partitions $\mu \subseteq \lambda$ with $\ell(\mu) = \ell(\lambda)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip.

Proof. Expand $GQ_\mu(x)$ in $\sum_{\mu} GQ_\mu(x)gp_\mu(y) = \sum_{\lambda} GP_\lambda(x)gq_\lambda(y)$ using Theorem 1.1 and then equate the coefficients of $GP_\lambda(x)$. \square

For example, $gq_{(m,\dots,3,2,1)} = 2^m gp_{(m,\dots,3,2,1)}$ for any $m \in \mathbb{N}$ and $gq_{(n)} = 2gp_{(n)} + \beta gp_{(n-1)}$ when $n \geq 2$. There is also an analogue of Corollary 1.2:

COROLLARY 6.3. *If λ is a strict partition with m parts then gq_λ is an $\mathbb{N}[\beta]$ -linear combination of gp -functions if and only if $\lambda - (m, \dots, 3, 2, 1)$ is also strict.*

Proof. One can check that if $\lambda - (m, \dots, 3, 2, 1)$ is strict then every μ indexing the sum in Corollary 6.2 has $\text{cols}(\lambda/\mu) = |\lambda/\mu|$, while if $\lambda - (m, \dots, 3, 2, 1)$ is not strict then at least one such μ has $\text{cols}(\lambda/\mu) = 1$ and $|\lambda/\mu| = 2$. \square

Recall that $\widehat{a}_{\mu\nu}^\lambda$ and $\widehat{b}_{\mu\nu}^\lambda$ are the integers appearing in the respective expansions of $GQ_{\lambda//\mu}$ and $GP_{\lambda//\mu}$ in (5.6). These numbers are zero if $|\mu| + |\nu| < |\lambda|$.

PROPOSITION 6.4. *If μ and ν are strict partitions then*

$$(6.2) \quad gp_\mu gp_\nu = \sum_\lambda \widehat{a}_{\mu\nu}^\lambda \beta^{|\mu|+|\nu|-|\lambda|} gp_\lambda \text{ and } gq_\mu gq_\nu = \sum_\lambda \widehat{b}_{\mu\nu}^\lambda \beta^{|\mu|+|\nu|-|\lambda|} gq_\lambda$$

where the sums are over all strict partitions λ .

Proof. This follows by a standard argument similar to the proof of [22, Prop. 8.2], which is equivalent to (6.2) when $\beta = 0$. Let $\Delta(x; y) := \prod_{i,j \geq 1} \frac{1-x_i y_j}{1-x_i y_j}$. Introduce a third set of commuting variables z_1, z_2, z_3, \dots . Then

$$\begin{aligned} \sum_\lambda \sum_\mu GQ_\mu(x) GQ_{\lambda//\mu}(y) gp_\lambda(z) &= \sum_\lambda GQ_\lambda(x, y) gp_\lambda(z) = \Delta(x, z) \Delta(y, z) \\ &= \left(\sum_\mu GQ_\mu(x) gp_\mu(z) \right) \left(\sum_\nu GQ_\nu(y) gp_\nu(z) \right) \end{aligned}$$

by equations (5.5) and (6.1). The first identity follows by substituting the formula (5.6) for $GQ_{\lambda//\mu}$ into the first expression and extracting the coefficients of $GQ_\mu(x) GQ_\nu(y)$. The second identity follows similarly. \square

Since $GP_{\lambda//\mu} = GQ_{\lambda//\mu} = 0$ when $\mu \not\subseteq \lambda$, we already know from (5.6) that

$$(6.3) \quad \widehat{a}_{\mu\nu}^\lambda = \widehat{b}_{\mu\nu}^\lambda = 0$$

when $\mu \not\subseteq \lambda$ (and, as noted above, when $|\lambda| > |\mu| + |\nu|$). We also have

$$(6.4) \quad \widehat{a}_{\mu\nu}^\lambda = \widehat{a}_{\nu\mu}^\lambda \quad \text{and} \quad \widehat{b}_{\mu\nu}^\lambda = \widehat{b}_{\nu\mu}^\lambda$$

by Proposition 6.4 since $\mathbb{Z}[\beta][[x_1, x_2, \dots]]$ is a commutative ring.

If μ and ν are strict partitions then [8, Props. 3.4 and 3.5] imply that

$$(6.5) \quad \begin{aligned} GP_\mu GP_\nu &= \sum_\lambda a_{\mu\nu}^\lambda \beta^{|\lambda|-|\mu|-|\nu|} GP_\lambda, \\ GQ_\mu GQ_\nu &= \sum_\lambda b_{\mu\nu}^\lambda \beta^{|\lambda|-|\mu|-|\nu|} GQ_\lambda, \end{aligned}$$

for unique integers $a_{\mu\nu}^\lambda, b_{\mu\nu}^\lambda \in \mathbb{Z}$. It is known from [3] that the GP -expansion is finite with every $a_{\mu\nu}^\lambda \in \mathbb{N}$, but *a priori* the coefficients $b_{\mu\nu}^\lambda$ could be nonzero for infinitely many strict partitions λ . We will discuss the problem of showing that the GQ -expansion is finite in Section 7. It is again clear that

$$(6.6) \quad a_{\mu\nu}^\lambda = a_{\nu\mu}^\lambda \quad \text{and} \quad b_{\mu\nu}^\lambda = b_{\nu\mu}^\lambda$$

but the following property requires an argument.

PROPOSITION 6.5. *One has $a_{\mu\nu}^\lambda = b_{\mu\nu}^\lambda = 0$ if $\mu \not\subseteq \lambda$ or $\nu \not\subseteq \lambda$ or $|\mu| + |\nu| > |\lambda|$.*

Proof. [8, Prop. 3.4] asserts that any finite product of GP -functions is a (possibly infinite) $\mathbb{Z}[\beta]$ -linear combination of GP -functions, while [8, Prop. 3.5] states an analogous property for the GQ -functions. Since GP_λ and GQ_λ are both homogeneous of degree $|\lambda|$ if we set $\deg(\beta) = -1$ and $\deg(x_i) = 1$, it must hold that $a_{\mu\nu}^\lambda = b_{\mu\nu}^\lambda = 0$ when $|\mu| + |\nu| > |\lambda|$.

It remains to show that both coefficients are zero when $\mu \not\subseteq \lambda$. This follows for $a_{\mu\nu}^\lambda$ from the Littlewood–Richardson rule for the GP_λ 's due to Clifford, Thomas, and Yong [3, Thm. 1.2]: in this result, the coefficient $C_{\mu\nu}^\lambda$ is equal to $a_{\mu\nu}^\lambda$ by [8, Cor. 8.1].

We turn to the $b_{\mu\nu}^\lambda$ coefficients. When p is a positive integer, there is a Pieri-type formula for the GQ -expansion of $GQ_\mu GQ_{(p)}$ due to Buch and Ravikumar [2, Cor. 5.6]: in this result, the coefficient $c_{\mu,p}^\lambda$ is equal to $b_{\mu\nu}^\lambda$ for $\nu = (p)$ again by [8, Cor. 8.1]. Buch and Ravikumar's formula implies that $b_{\mu\nu}^\lambda = 0$ if $\mu \not\subseteq \lambda$ and $\ell(\nu) \leq 1$. It follows that if p_1, p_2, \dots, p_k are any positive integers then the product $GQ_\mu GQ_{(p_1)} GQ_{(p_2)} \cdots GQ_{(p_k)}$ is a possibly infinite $\mathbb{Z}[\beta]$ -linear combination of GQ_λ 's indexed by strict partitions with $\lambda \supseteq \mu$.

From this observation, to complete the proof it is enough to show that each GQ_ν is a possibly infinite $\mathbb{Z}[\beta]$ -linear combination of products of the form $GQ_{(p_1)} GQ_{(p_2)} \cdots GQ_{(p_k)}$. This claim is a consequence of [17, Thm. 5.8], which gives a formula for a more general *universal factorial Hall–Littlewood Q -function* $HQ_\nu^L(\mathbf{x}_n; t \mid \mathbf{b})$, essentially as a linear combination of products of other such functions indexed by one-row partitions. The notation in [17] makes it slightly nontrivial to connect this formula to our situation. However, the property needed for GQ_ν follows directly from the discussion in [17, §5.2.4] (in particular, from [17, Thm. 5.10]), once one observes that the generating function

$$GQ(u \mid \mathbf{b})^{(k)} := \frac{1}{1 + \beta u} \prod_{j=1}^n \frac{1 + (u^{-1} + \beta)x_j}{1 + (u^{-1} + \beta)\bar{x}_j} \times \prod_{j=1}^k (1 + (u^{-1} + \beta)b_j)$$

in [17, Eq. (5.10)] reduces when $\mathbf{b} = \mathbf{0}$ and $n \rightarrow \infty$ to the expression below:

LEMMA 6.6. *When expanded as a power series in u^{-1} , the expression*

$$\frac{1}{1 + \beta u} \prod_{j=1}^{\infty} \frac{1 + (u^{-1} + \beta)x_j}{1 + (u^{-1} + \beta)\bar{x}_j}$$

is equal to $\sum_{n \in \mathbb{Z}} GQ_{(n)} u^{-n}$ where we set $GQ_{(n)} := (-\beta)^{-n}$ for $n < 0$.

Proof. This is essentially [7, Rem. 5.11] given [7, Def. 3.5]. Here is a self-contained proof explained to us by Joel Lewis. Let $\ell(T)$ denote the number boxes in a tableau T . Then $\sum_{n > 0} GQ_{(n)} t^n = \sum_T \beta^{|T| - \ell(T)} x^T t^{\ell(T)}$ where the sum is over all semistandard set-valued shifted tableaux T with a nonempty one-row shape. Such a tableau T is specified uniquely by the following choices:

- The finite set of positive integers $\{j_1 < \dots < j_k\}$ such that at least one of j_i or j'_i appears in some box of T .
- The numbers $n_1, \dots, n_k > 0$ such that j_i or j'_i appears in n_i boxes of T .
- For each $i \in [k]$, whether the first box containing j_i or j'_i contains just j_i , just j'_i , or both j_i and j'_i .
- For each $i > 1$, whether the first box containing j_i or j'_i contains no smaller numbers, or contains at least one of j_{i-1} or j'_{i-1} .

For the tableau T corresponding to this data, the value of $|T|$ (respectively, $\ell(T)$) depends on the numbers n_1, \dots, n_k and the choices in the third (respectively, fourth)

bullet point. It follows that we can write $\sum_{n>0} GQ_{(n)}t^n$ as

$$\sum_{\substack{k>0 \\ 0<j_1<\dots<j_k \\ n_1,\dots,n_k>0}} (2x_{j_1} + \beta x_{j_1}^2)t \times (x_{j_1}t)^{n_1-1} \times \prod_{i=2}^k (2\beta x_{j_i} + \beta^2 x_{j_i}^2) \left(\frac{t}{\beta} + 1\right) \times (x_{j_i}t)^{n_i-1}$$

Fixing k and $j_1 < \dots < j_k$ and summing over n_1, \dots, n_k turns this into

$$\sum_{\substack{k>0 \\ 0<j_1<\dots<j_k}} \frac{(2x_{j_1} + \beta x_{j_1}^2)t}{1 - x_{j_1}t} \times \prod_{i=2}^k \frac{(2x_{j_i} + \beta x_{j_i}^2)(t + \beta)}{1 - x_{j_i}t}.$$

Pulling out a factor of $\frac{t}{t+\beta} = \frac{1}{1+\beta t^{-1}}$ transforms this to

$$\sum_{\substack{S \subseteq \{1,2,\dots\} \\ 0 < |S| < \infty}} \frac{t}{t+\beta} \prod_{j \in S} \frac{(2x_j + \beta x_j^2)(t+\beta)}{1 - x_j t} = \frac{1}{1+\beta t^{-1}} \left(\prod_{j=1}^{\infty} \left(1 + \frac{(2x_j + \beta x_j^2)(t+\beta)}{1 - x_j t} \right) - 1 \right).$$

Since $1 + \frac{(2x+\beta x^2)(t+\beta)}{1-xt} = \frac{(1+\beta x)(1+\beta x+xt)}{1-xt} = \frac{1+(t+\beta)x}{1+(t+\beta)x}$ we conclude that

$$\sum_{n>0} GQ_{(n)}t^n = \frac{1}{1 + \beta t^{-1}} \prod_{j=1}^{\infty} \frac{1 + (t + \beta)x_j}{1 + (t + \beta)x_j} - \frac{1}{1 + \beta t^{-1}}.$$

The desired formula follows by replacing the formal parameter t by u^{-1} . □

This completes the proof of Proposition 6.5. □

For any strict partitions λ and μ we define

$$(6.7) \quad gq_{\lambda/\mu} := \sum_{\nu} a_{\mu\nu}^{\lambda} \beta^{|\lambda|-|\mu|-|\nu|} gq_{\nu} \quad \text{and} \quad gp_{\lambda/\mu} := \sum_{\nu} b_{\mu\nu}^{\lambda} \beta^{|\lambda|-|\mu|-|\nu|} gp_{\nu}$$

where ν ranges over all strict partitions. Proposition 6.5 implies that these symmetric functions have bounded degree and are zero whenever $\mu \not\subseteq \lambda$. As gp_{λ} and gq_{λ} are homogeneous of degree $|\lambda|$ when $\deg(\beta) = 1$, it follows that $gp_{\lambda/\mu}$ and $gq_{\lambda/\mu}$ are homogeneous of degree $|\lambda| - |\mu|$ when nonzero.

PROPOSITION 6.7. *If λ is a strict partition then*

$$(6.8) \quad gp_{\lambda}(x, y) = \sum_{\nu} gp_{\nu}(x)gp_{\lambda/\nu}(y) \quad \text{and} \quad gq_{\lambda}(x, y) = \sum_{\nu} gq_{\nu}(x)gq_{\lambda/\nu}(y)$$

where both sums are over all strict partitions ν .

Proof. These identities are equivalent to the first two parts of [16, Prop. 3.2]. We show how to derive the first identity for completeness. Observe that

$$\begin{aligned} \sum_{\lambda} GQ_{\lambda}(x)gp_{\lambda}(y, z) &= \Delta(x, y)\Delta(x, z) \\ &= \left(\sum_{\mu} GQ_{\mu}(x)gp_{\mu}(y) \right) \left(\sum_{\nu} GQ_{\nu}(x)gp_{\nu}(z) \right) \\ &= \sum_{\mu, \nu, \lambda} b_{\mu\nu}^{\lambda} \beta^{|\lambda|-|\mu|-|\nu|} GQ_{\lambda}(x)gp_{\mu}(y)gq_{\nu}(z) \\ &= \sum_{\lambda} \left(GQ_{\lambda}(x) \sum_{\mu} gp_{\mu}(y)gp_{\lambda/\mu}(z) \right) \end{aligned}$$

by (6.1) and (6.7). Now equate the coefficients of $GQ_{\lambda}(x)$. □

REMARK 6.8. Setting $\beta = 0$ transforms (6.8) to (5.2), so $P_{\lambda/\mu} = gp_{\lambda/\mu}|_{\beta=0}$ and $Q_{\lambda/\mu} = gq_{\lambda/\mu}|_{\beta=0}$. Thus $gp_{\lambda/\mu}$ and $gq_{\lambda/\mu}$ are each nonzero if and only if $\mu \subseteq \lambda$.

We may now prove a dual version of Theorem 5.3.

THEOREM 6.9. *Suppose $\kappa \subseteq \lambda$ are strict partitions. Then*

$$gq_{\lambda/\kappa} = \sum_{(\nu, \mu)} 2^{\ell(\lambda) - \ell(\kappa) + \text{overlap}(\nu/\kappa)} (-1)^{\text{cols}(\lambda/\mu)} (-\beta/2)^{|\nu/\kappa| + |\lambda/\mu|} gp_{\mu/\nu}$$

where the sum is over all pairs of strict partitions (ν, μ) with $\kappa \subseteq \nu \subseteq \mu \subseteq \lambda$ and $\ell(\kappa) = \ell(\nu) \leq \ell(\mu) = \ell(\lambda)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip.

REMARK 6.10. If there were a bilinear form that made $\{gq_{\lambda/\kappa}\}$ the dual basis of $\{GP_{\lambda/\kappa}\}$ and $\{gp_{\mu/\nu}\}$ the dual basis of $\{GQ_{\mu/\nu}\}$, then this result would follow immediately from Theorem 5.3. Since the various skew functions are not even linearly independent, no such form is available and we need another argument.

Proof. Expanding $gq_{\lambda}(x, y)$ using Corollary 6.2 and then applying (6.8) gives

$$gq_{\lambda}(x, y) = \sum_{\mu} \sum_{\kappa} 2^{\ell(\lambda)} (-1)^{\text{cols}(\lambda/\mu)} (-\beta/2)^{|\lambda/\mu|} gp_{\kappa}(x) gp_{\mu/\kappa}(y)$$

where the outer sum is over all strict partitions μ with $\mu \subseteq \lambda$ and $\ell(\mu) = \ell(\lambda)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip and the inner sum is over strict partitions κ with $\kappa \subseteq \mu$ (since $gp_{\mu/\kappa} = 0$ if $\kappa \not\subseteq \mu$). Alternatively, using Corollary 6.2 to expand the right side of (6.8) gives

$$gq_{\lambda}(x, y) = \sum_{\eta} \sum_{\kappa} 2^{\ell(\kappa)} (-1)^{\text{cols}(\eta/\kappa)} (-\beta/2)^{|\eta/\kappa|} gp_{\kappa}(x) gq_{\lambda/\eta}(y)$$

where the outer sum is over all strict partitions η with $\eta \subseteq \lambda$ (since $gq_{\lambda/\eta} = 0$ when $\eta \not\subseteq \lambda$) and the inner sum over κ with $\kappa \subseteq \eta$ and $\ell(\kappa) = \ell(\eta)$ such that $\text{SD}_{\eta/\kappa}$ is a vertical strip. Equating coefficients of gp_{κ} gives

$$\sum_{\eta} 2^{\ell(\kappa)} (-1)^{\text{cols}(\eta/\kappa)} \left(\frac{-\beta}{2}\right)^{|\eta/\kappa|} gq_{\lambda/\eta} = \sum_{\mu} 2^{\ell(\lambda)} (-1)^{\text{cols}(\lambda/\mu)} \left(\frac{-\beta}{2}\right)^{|\lambda/\mu|} gp_{\mu/\kappa}$$

where the sums are over certain strict partitions η and μ ; we can rewrite this as

$$gq_{\lambda/\kappa} = \sum_{\mu} 2^{\ell(\lambda) - \ell(\kappa)} (-1)^{\text{cols}(\lambda/\mu)} \left(\frac{-\beta}{2}\right)^{|\lambda/\mu|} gp_{\mu/\kappa} - \sum_{\eta} (-1)^{\text{cols}(\eta/\kappa)} \left(\frac{-\beta}{2}\right)^{|\eta/\kappa|} gq_{\lambda/\eta}$$

where the first sum is over all strict partitions μ with $\kappa \subseteq \mu \subseteq \lambda$ and $\ell(\mu) = \ell(\lambda)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip, and the second sum is over all strict partitions η with $\kappa \subsetneq \eta \subseteq \lambda$ and $\ell(\kappa) = \ell(\eta)$ such that $\text{SD}_{\eta/\kappa}$ is a vertical strip.

If $\kappa = \lambda$ then $gq_{\lambda/\kappa} = 1 = gp_{\lambda/\kappa}$ as the theorem predicts. Otherwise, we may assume by induction that the desired formula holds for $gq_{\lambda/\eta}$ with $\kappa \subsetneq \eta \subseteq \lambda$. Substituting these formulas into the equation above expands $gq_{\lambda/\kappa}$ as a linear combination of $gp_{\mu/\nu}$'s where μ and ν range over the strict partitions with $\kappa \subseteq \nu \subseteq \mu \subseteq \lambda$ and $\ell(\kappa) = \ell(\nu) \leq \ell(\mu) = \ell(\lambda)$ such that $\text{SD}_{\lambda/\mu}$ is a vertical strip. The coefficient of $gp_{\mu/\kappa}$ in this expansion is $2^{\ell(\lambda) - \ell(\kappa)} (-1)^{\text{cols}(\lambda/\mu)} \left(\frac{-\beta}{2}\right)^{|\lambda/\mu|}$ as desired. The coefficient of $gp_{\mu/\nu}$ when $\kappa \subsetneq \nu \subseteq \mu \subseteq \lambda$ and $\ell(\kappa) = \ell(\nu) \leq \ell(\mu) = \ell(\lambda)$ and $\text{SD}_{\lambda/\mu}$ is a vertical strip is the sum

$$- \sum_{\eta} (-1)^{\text{cols}(\eta/\kappa)} \left(\frac{-\beta}{2}\right)^{|\eta/\kappa|} \left(2^{\ell(\lambda) - \ell(\eta) + \text{overlap}(\nu/\eta)} (-1)^{\text{cols}(\lambda/\mu)} \left(\frac{-\beta}{2}\right)^{|\nu/\eta| + |\lambda/\mu|} \right)$$

over all strict partitions η where $\kappa \subsetneq \eta \subseteq \nu$ and $\text{SD}_{\eta/\kappa}$ is a vertical strip. Rewriting this as $2^{\ell(\lambda) - \ell(\eta)} (-1)^{\text{cols}(\lambda/\mu)} \left(\frac{-\beta}{2}\right)^{|\lambda/\mu| + |\nu/\kappa|} \sum_{\eta} (-1)^{\text{cols}(\eta/\kappa) + 1} 2^{\text{overlap}(\nu/\eta)}$, we see that it suffices to show that $\sum_{\eta} (-1)^{\text{cols}(\eta/\kappa) + 1} 2^{\text{overlap}(\nu/\eta)} = 2^{\text{overlap}(\nu/\kappa)}$ where the sum is

again over η with $\kappa \subsetneq \eta \subseteq \nu$ such that $SD_{\eta/\kappa}$ is a vertical strip. After rearranging terms, this identity is the second half of Lemma 5.1. \square

By putting everything together we can also prove a generalization of (6.1). This gives two shifted analogues of Yeliussizov’s skew Cauchy identity for *symmetric Grothendieck polynomials* [24, Thm. 5.1].

THEOREM 6.11. *Let μ and ν be strict partitions. Then*

$$(6.9) \quad \sum_{\lambda} GP_{\lambda//\mu}(x)gq_{\lambda/\nu}(y) = \prod_{i,j \geq 1} \frac{1 - \bar{x}_i y_j}{1 - x_i y_j} \sum_{\kappa} GP_{\nu//\kappa}(x)gq_{\mu/\kappa}(y)$$

and

$$(6.10) \quad \sum_{\lambda} GQ_{\lambda//\mu}(x)gp_{\lambda/\nu}(y) = \prod_{i,j \geq 1} \frac{1 - \bar{x}_i y_j}{1 - x_i y_j} \sum_{\kappa} GQ_{\nu//\kappa}(x)gp_{\mu/\kappa}(y)$$

where $\bar{x}_i := \frac{-x_i}{1+\beta x_i}$ and where λ and κ range over all strict partitions.⁽²⁾

Proof. Let w_1, w_2, w_3, \dots be a fourth set of commuting variables. The first expression in (6.9) is the coefficient of $GP_{\mu}(w)gq_{\nu}(z)$ in

$$\sum_{\lambda} GP_{\lambda}(w, x)gq_{\lambda}(z, y) = \Delta(w; z)\Delta(w; y)\Delta(x; z)\Delta(x; y)$$

by (5.5), (6.1), and (6.8). Since $\Delta(x; y) = \prod_{i,j \geq 1} \frac{1 - \bar{x}_i y_j}{1 - x_i y_j}$, to prove (6.9) it suffices to show that $\sum_{\kappa} GP_{\nu//\kappa}(x)gq_{\mu/\kappa}(y)$ is the coefficient of $GP_{\mu}(w)gq_{\nu}(z)$ in $\Delta(w; z)\Delta(w; y)\Delta(x; z)$. By (6.1) this product is equal to

$$\sum_{\kappa} \sum_{\gamma} \sum_{\lambda} GP_{\kappa}(w)gq_{\kappa}(z) \cdot GP_{\gamma}(w)gq_{\gamma}(y) \cdot GP_{\lambda}(x)gq_{\lambda}(z).$$

Once we rearrange the terms of this expression as

$$\sum_{\kappa} \sum_{\gamma} \sum_{\lambda} GP_{\kappa}(w)GP_{\gamma}(w) \cdot GP_{\lambda}(x)gq_{\gamma}(y) \cdot gq_{\kappa}(z)gq_{\lambda}(z)$$

it follows by (6.2) and (6.5) that it is equal to

$$\sum_{\kappa} \sum_{\gamma} \sum_{\lambda} \sum_{\mu} \sum_{\nu} \beta^{|\mu|-|\kappa|-|\gamma|} a_{\kappa\gamma}^{\mu} GP_{\mu}(w) \cdot GP_{\lambda}(x)gq_{\gamma}(y) \cdot \beta^{|\kappa|+|\lambda|-|\nu|} \widehat{b}_{\kappa\lambda}^{\nu} gq_{\nu}(z).$$

On rearranging the terms of this expression to be

$$\sum_{\kappa} \sum_{\gamma} \sum_{\lambda} \sum_{\mu} \sum_{\nu} GP_{\mu}(w) \cdot \beta^{|\kappa|+|\lambda|-|\nu|} \widehat{b}_{\kappa\lambda}^{\nu} GP_{\lambda}(x) \cdot \beta^{|\mu|-|\kappa|-|\gamma|} a_{\kappa\gamma}^{\mu} gq_{\gamma}(y) \cdot gq_{\nu}(z)$$

we deduce by (5.6) and (6.7) that

$$\Delta(w; z)\Delta(w; y)\Delta(x; z) = \sum_{\kappa} \sum_{\mu} \sum_{\nu} GP_{\mu}(w) \cdot GP_{\nu//\kappa}(x) \cdot gq_{\mu/\kappa}(y) \cdot gq_{\nu}(z).$$

The coefficient of $GP_{\mu}(w)gq_{\nu}(z)$ in the last expression is $\sum_{\kappa} GP_{\nu//\kappa}(x)gq_{\mu/\kappa}(y)$, so (6.9) holds. The second identity (6.10) follows by the same argument after interchanging the symbols $GP \leftrightarrow GQ$, $gp \leftrightarrow gq$, $a \leftrightarrow b$ and $\widehat{a} \leftrightarrow \widehat{b}$. \square

⁽²⁾Equivalently, one can restrict λ to the strict partitions containing both μ and ν since otherwise $GQ_{\lambda//\mu}(x)gp_{\lambda/\nu}(y) = 0$, and one can restrict κ to the strict partitions contained in both μ and ν since otherwise $GQ_{\nu//\kappa}(x)gp_{\mu/\kappa} = 0$.

We could then derive $gp_{\lambda/\nu} = \sum_{T \in \text{ShRPP}_P(\lambda/\nu)} (-\beta)^{|\lambda/\nu| - |\text{wtRPP}(T)|} x^{\text{wtRPP}(T)}$ by comparing the preceding identity with (6.8) and equating the coefficients of gp_ν . The reductions in gq -case are similar.

Nakagawa and Naruse give another conjectural formula for gq_λ as a certain Pfaffian in [17, §6.2], but we will not discuss this here.

Write λ^\top for the transpose of a partition λ . Let ω be the operator acting on (possibly infinite) $\mathbb{Z}[\beta]$ -linear combinations of Schur functions $f = \sum_\lambda c_\lambda s_\lambda$ as $\omega(f) := \sum_\lambda c_\lambda s_{\lambda^\top}$. For strict partitions μ and λ define

$$(7.2) \quad jp_{\lambda/\mu} := \omega(gp_{\lambda/\mu}) \quad \text{and} \quad jq_{\lambda/\mu} := \omega(gq_{\lambda/\mu}),$$

setting $jp_\lambda := jp_{\lambda/\emptyset}$ and $jq_\lambda := jq_{\lambda/\emptyset}$. These are finite $\mathbb{Z}[\beta]$ -linear combinations of Schur functions which are nonzero if and only if $\mu \subseteq \lambda$, since the same is true of the gp - and gq -functions. When nonzero, $jp_{\lambda/\mu}$ and $jq_{\lambda/\mu}$ are homogeneous of degree $|\lambda| - |\mu|$ under the convention that $\deg(\beta) = \deg(x_i) = 1$.

As ω fixes all skew Schur P - and Q -functions [12, Ex. 3(a), §III.8], we have

$$(7.3) \quad P_{\lambda/\mu} = jp_{\lambda/\mu}|_{\beta=0} \quad \text{and} \quad Q_{\lambda/\mu} = jq_{\lambda/\mu}|_{\beta=0}.$$

Additionally, since ω is an automorphism of the Hopf algebra of bounded degree symmetric power series in $\mathbb{Z}[\beta][[x_1, x_2, \dots]]$, the identity (6.8) is equivalent to

$$(7.4) \quad jp_\lambda(x, y) = \sum_\mu jp_\mu(x) jp_{\lambda/\mu}(y) \quad \text{and} \quad jq_\lambda(x, y) = \sum_\mu jq_\mu(x) jq_{\lambda/\mu}(y)$$

where both sums are over all strict partitions μ .

We may also describe a conjectural generating function formula for $jp_{\lambda/\mu}$ and $jq_{\lambda/\mu}$. This formula appears to be new.

A partition of a set S is a set Π of disjoint nonempty subsets $B \subseteq S$, called *blocks*, with $S = \bigsqcup_{B \in \Pi} B$. Assume $\mu \subseteq \lambda$. We define a *shifted bar tableau* of shape λ/μ to be a pair $T = (V, \Pi)$, where V is a semistandard shifted tableau of shape λ/μ and Π is a partition of $\text{SD}_{\lambda/\mu}$ such that each block $B \in \Pi$ is a set of adjacent boxes containing the same entry in V . Because V is semistandard, each of these blocks must consist of a contiguous “bar” within a single row or single column. One might draw a shifted bar tableau as a picture like

$$(7.5) \quad \begin{array}{cccc} & 2 & & 3' \\ 1 & & 1 & & 3 \end{array}$$

to represent

$$T = (V, \Pi) = \left(\left(\begin{array}{|c|c|c|c|} \hline 2 & 2 & 3' & \\ \hline 1 & 1 & 1 & 3' & 3 \\ \hline \end{array} \right), \left(\begin{array}{|c|c|c|c|} \hline \cdot & \cdot & \cdot & \\ \hline \cdot & \cdot & \cdot & \cdot & \cdot \\ \hline \end{array} \right) \right).$$

Let $\text{ShBT}_Q(\lambda/\mu)$ denote the set of all shifted bar tableaux of shape λ/μ and let $\text{ShBT}_P(\lambda/\mu)$ be the subset of such pairs $T = (V, \Pi)$ where V has no primed diagonal entries. Given $T = (V, \Pi) \in \text{ShBT}_Q(\lambda/\mu)$ we set

$$|T| := |\Pi| \quad \text{and} \quad x^T := \prod_{i \geq 1} x_i^{b_i}$$

where b_i is the number of blocks in Π containing i or i' in V . The example T shown in (7.5) belongs to $\text{ShBT}_P(\lambda/\mu)$ for $\lambda = (5, 3)$ and $\mu = \emptyset$ and has $|T| = 5$ and $x^T = x_1^2 x_2 x_3^2$.

If every block in Π has size one then $|T| = |\lambda/\mu|$ and $x^T = x^V$. If Π has as few blocks as possible given V , then $x^T = x^{\text{wtRPP}(V)}$. Finally, observe that if V is fixed, then the sum of x^T over all Π such that $T = (V, \Pi)$ is a shifted bar tableau is

$\prod_{i \geq 1} x_i^{r_i + c_i} (x_i + 1)^{m_i - r_i - c_i}$ where r_i is the number of rows of V containing an entry equal to i , c_i is the number of columns of V containing an entry equal to i' , and m_i is the number of boxes of V containing i or i' .


CONJECTURE 7.4. *If $\mu \subseteq \lambda$ are strict partitions then*

$$jp_{\lambda/\mu} = \sum_{T \in \text{ShBT}_P(\lambda/\mu)} (-\beta)^{|\lambda/\mu| - |T|} x^T,$$

$$jq_{\lambda/\mu} = \sum_{T \in \text{ShBT}_Q(\lambda/\mu)} (-\beta)^{|\lambda/\mu| - |T|} x^T.$$

As with Conjecture 7.1, to prove this result it would suffice to assume $\mu = \emptyset$.


EXAMPLE 7.5. If $\lambda = (2, 1)$ and $\mu = \emptyset$ then $\text{ShBT}_P(\lambda/\mu)$ consists of



for all positive integers $a < b < c$, so Conjecture 7.4 asserts that

$$jp_{21} = 2 \sum_{a < b < c} x_a x_b x_c + \sum_{a < b} (x_a^2 x_b + x_a x_b^2) - \beta \sum_{a < b} x_a x_b = s_{21} - \beta s_{11}.$$

As in Example 7.2, there is a weight-preserving 4-to-1 map $\text{ShBT}_Q(21/\emptyset) \rightarrow \text{ShBT}_P(21/\emptyset)$; this is given by either removing all diagonal primes or applying



Thus Conjecture 7.4 also predicts that $jq_{21} = 4s_{21} - 4\beta s_{11}$. These formulas are consistent with Example 7.2 as $jp_{21} = \omega(gp_{21})$ and $jq_{21} = \omega(gq_{21})$.

REMARK 7.6. One can systematically test Conjecture 7.1 by substituting into the Cauchy identity (6.1) both the set-valued generating functions for GP_λ and GQ_λ and the predicted reverse plane partition generating functions for gp_λ and gq_λ , then truncating all three expressions in (6.1) to finitely many variables and finite degree, and finally checking that the resulting polynomials match. All computer calculations we have done along these lines support the conjecture.

To test Conjecture 7.4, one can compute the Schur polynomial expansions of the right hand expressions in Conjecture 7.1 and 7.4 restricted to finitely many variables x_1, x_2, \dots, x_n . If n is large enough, then the corresponding expansions should be related by transposing all partition indices. We have used a computer to verify Conjecture 7.4 in this way for all strict partitions λ with $|\lambda| \leq 6$.

Conjectures 7.1 and 7.4 are shifted analogues of results in [10, §9]. One may be able to adapt the operator methods used there and in [24] to prove both formulas. We will not pursue this here, beyond verifying the one-row case:

PROPOSITION 7.7. *Conjectures 7.1 and 7.4 hold when $\ell(\lambda) \leq 1$.*

Proof. When $\ell(\lambda) = 0$ the conjectures assert that $gp_\emptyset = gq_\emptyset = jp_\emptyset = jq_\emptyset = 1$. This follows by comparing constant coefficients in (6.1) since $GP_\emptyset = GQ_\emptyset = 1$.

Suppose $\ell(\lambda) = 1$. We may assume $\mu = \emptyset$ in view of (6.8) and (7.4). We may also assume $\beta = -1$ since if f is $gp_\lambda, gq_\lambda, jp_\lambda, jq_\lambda$, or one of the conjectural generating functions, then by Proposition 6.1 we can recover f from $f|_{\beta=-1}$ by substituting $x_i \mapsto -\beta^{-1}x_i$ for all i and then multiplying the result by $(-\beta)^{|\lambda|}$.

Let $\overline{gp}_\lambda := \sum_{T \in \text{ShRPP}_P(\lambda)} x^{\text{wt}_{\text{RPP}}(T)}$ and $\overline{gq}_\lambda := \sum_{T \in \text{ShRPP}_Q(\lambda)} x^{\text{wt}_{\text{RPP}}(T)}$. We wish to show that $gp_{(n)} = \overline{gp}_{(n)}$ and $gq_{(n)} = \overline{gq}_{(n)}$ when n is a positive integer. Nakagawa

and Naruse assert that these identities hold [16, Prop. 5.2] but do not provide a proof. Here is an argument. The function $GP_\lambda(t)$ obtained by setting $x_1 = t$ and $x_2 = x_3 = \dots = 0$ is t^n if $\lambda = (n)$ and 0 if $\ell(\lambda) > 1$. Thus, if we set $x_1 = t$ and $x_i = 0$ for $i > 1$ then (6.1) becomes

$$\sum_{n \geq 0} t^n gq_{(n)}(y) = \prod_{j \geq 1} \frac{1 - \frac{-t}{1-t} y_j}{1 - t y_j}.$$

Rewriting y_j as x_j , we deduce that $gq_{(n)} = gq_{(n)}(x)$ is the coefficient of t^n in

$$\prod_{j \geq 1} \frac{1 - \frac{-t}{1-t} x_j}{1 - t x_j} = \prod_{j \geq 1} (1 + x_j t + x_j t^2 + x_j t^3 + \dots)(1 + x_j t + x_j^2 t^2 + x_j^3 t^3 + \dots).$$

Each way of expanding this product into monomials corresponds to a unique one-row shifted reverse plane partition T : if we multiply the a th term of $(1 + x_j t + x_j t^2 + x_j t^3 + \dots)$ with the b th term of $(1 + x_j t + x_j^2 t^2 + x_j^3 t^3 + \dots)$, considering 1 to be the 0th term of both sums, then the resulting monomial is $x^{\text{wt}_{\text{RPP}}(T)} t^{|T|}$ where T is the one-row shifted reverse plane partition with exactly $a = a(j)$ entries equal to j' and $b = b(j)$ entries equal to j for each $j \geq 1$. It follows that the coefficient of t^n in the above product is also $\overline{gq}_{(n)}$, so $gq_{(n)} = \overline{gq}_{(n)}$.

We have $\overline{gq}_{(1)} = 2\overline{gp}_{(1)} = 2(x_1 + x_2 + x_3 + \dots)$, and if $n \geq 2$ then $\overline{gp}_{(n)} - \overline{gp}_{(n-1)} = \sum_T x^{\text{wt}_{\text{RPP}}(T)}$ where the sum is over one-row shifted reverse plane partitions T of size n whose first two boxes contain distinct entries, the first of which is primed. Removing the prime from the first entry defines a weight-preserving bijection from the set of such T to $\text{ShRPP}_Q(n) - \text{ShRPP}_P(n)$, so it follows that $\overline{gq}_{(n)} = 2\overline{gp}_{(n)} - \overline{gp}_{(n-1)}$ when $n \geq 2$. Since we likewise have $gq_{(1)} = 2gp_{(1)}$ and $gq_{(n)} = 2gp_{(n)} - gp_{(n-1)}$ when $n \geq 2$ by Corollary 6.2, we deduce by induction that $gp_{(n)} = \overline{gp}_{(n)}$ for all n .

An (*unshifted*) *reverse plane partition* of shape λ is a filling of the (*unshifted*) diagram $D_\lambda := \{(i, j) : 1 \leq j \leq \lambda_i\}$ by positive integers, such that rows and columns are weakly increasing. The weight of such an object is defined in the same way as for shifted reverse plane partitions. As noted in [16, Prop. 5.3], it is easy to see that $\overline{gp}_{(n)} = \sum_T x^{\text{wt}_{\text{RPP}}(T)}$ where the sum is over all reverse plane partitions T of any of the hook shapes $a1^{n-a} := (a, 1, 1, \dots, 1)$ for $a \in [n]$; the relevant weight-preserving bijection is given by moving all primed entries in an element of $\text{ShRPP}_P(n)$ from the first row to the first column with primes removed. This sum is precisely $\sum_{a=1}^n g_{a1^{n-a}}$ where g_λ is the *dual stable Grothendieck polynomial* discussed, for example, in [10, §9.1].

An (*unshifted*) *bar tableau* of shape λ is defined in the same way as a shifted bar tableau, except the underlying tableau is a filling of D_λ (rather than SD_λ) by positive integers (rather than positive half-integers); this is called a *valued-set tableau* in [10, §9.8]. The weight x^T is the same as in the shifted case. Let $\overline{jp}_\lambda := \sum_{T \in \text{ShBT}_P(\lambda)} x^T$ and $\overline{jq}_\lambda := \sum_{T \in \text{ShBT}_Q(\lambda)} x^T$. Then $\overline{jp}_{(n)} = \sum_T x^T$ where the sum is over all bar tableaux T of any of the hook shapes $a1^{n-a}$ for $a \in [n]$; the relevant weight-preserving bijection is again given by moving all primed entries in an element of $\text{ShBT}_P(n)$ from the first row to the first column with primes removed. (Each primed entry comprised its own block in the first row and is assigned to its own block in the first column.) This sum is precisely $\sum_{a=1}^n j_{a1^{n-a}}$ where j_λ is the generating function discussed in [10, §9.8], which has $j_\lambda = \omega(g_\lambda)$ for all partitions λ [10, Prop. 9.25].

Combining the last two paragraphs shows that $\overline{jp}_{(n)} = \omega(\overline{gp}_{(n)})$ so $\overline{jp}_{(n)} = \omega(gp_{(n)}) = \omega(\overline{gp}_{(n)}) = \overline{jp}_{(n)}$ for all n . Finally, we have $\overline{jq}_{(1)} = 2\overline{jp}_{(1)} = 2(x_1 + x_2 + x_3 + \dots)$, and if $n \geq 2$ then $\overline{jp}_{(n)} - \overline{jp}_{(n-1)} = \sum_T x^T$ where the sum is over one-row shifted bar tableaux T of size n whose first two entries are

unprimed but not in the same block. Adding a prime to the diagonal entry gives a weight-preserving bijection from the set of such T to $\text{ShBT}_Q(n) - \text{ShBT}_P(n)$, so $\overline{jq}_{(n)} = 2\overline{jp}_{(n)} - \overline{jp}_{(n-1)}$ when $n \geq 2$. Since the same formulas relate $jq_{(n)}$ to $jp_{(n)}$ by the linearity of ω , we must have $jq_{(n)} = \overline{jq}_{(n)}$. \square

Conjecture 7.4 has some consequences regarding the numbers $a_{\mu\nu}^\lambda$ and $b_{\mu\nu}^\lambda$.

THEOREM 7.8. *Suppose the formula for $jq_{\lambda/\mu}$ (respectively, $jp_{\lambda/\mu}$) in Conjecture 7.4 holds. Then the coefficients in the product expansions (6.5) satisfy $a_{\mu\nu}^\lambda = 0$ (respectively, $b_{\mu\nu}^\lambda = 0$) whenever $\ell(\lambda) > \ell(\mu) + \ell(\nu)$.*

Proof. Our argument is based on Yeliussizov’s proof of [24, Thm. 8.4]. Since consecutive diagonal entries in a semistandard shifted tableau differ by at least one, the assumed formula in Conjecture 7.4 implies that $jq_\nu(x_1, \dots, x_n) \neq 0$ if and only if $\ell(\nu) \leq n$ and that $jq_{\lambda/\mu}(x_1, \dots, x_n) = 0$ whenever $\ell(\lambda) > n + \ell(\mu)$. As $jq_\nu|_{\beta=0} = Q_\nu$, the set of polynomials $\{jq_\kappa(x_1, \dots, x_n) : \ell(\kappa) \leq n\}$ is linearly independent over $\mathbb{Z}[\beta]$ since $\{Q_\kappa(x_1, \dots, x_n) : \ell(\kappa) \leq n\}$ is linearly independent.

Applying ω to (6.7) gives $jq_{\lambda/\mu} = \sum_\kappa a_{\mu\kappa}^\lambda \beta^{|\lambda|-|\mu|-|\kappa|} jq_\kappa$. Thus if $a_{\mu\nu}^\lambda \neq 0$ and $n = \ell(\nu)$ then $jq_{\lambda/\mu}(x_1, \dots, x_n) = \sum_{\ell(\kappa) \leq n} a_{\mu\kappa}^\lambda \beta^{|\lambda|-|\mu|-|\kappa|} jq_\kappa(x_1, \dots, x_n) \neq 0$. But this means that $\ell(\lambda) \leq n + \ell(\mu) = \ell(\mu) + \ell(\nu)$ as desired. The claim about $b_{\mu\nu}^\lambda$ follows by the same argument after swapping $jq \leftrightarrow jp$ and $Q \leftrightarrow P$. \square

Ikeda and Naruse conjectured that the product expansions in (6.5) both have finitely many nonzero terms [8, Conj. 3.1 and 3.2]. For the GP -functions, this follows from results in [3], which also establish that each coefficient $a_{\mu\nu}^\lambda \in \mathbb{N}$; for other proofs see [6, §4], [13, §1.2], or [20, §8].

The same claim for the GQ -functions appears still to be open, but would be a consequence of Conjecture 7.4 by the following corollary. Specifically, this corollary shows that Conjecture 7.4 implies [8, Conj. 3.2]. Even given these conjectures, it is still an open problem to find a Littlewood–Richardson rule to compute $b_{\mu\nu}^\lambda$ outside the Pieri formula case $\nu = (p)$ handled in [2].

COROLLARY 7.9. *Suppose the formula for $jp_{\lambda/\mu}$ in Conjecture 7.4 holds. Then each product $GQ_\mu GQ_\nu$ is a finite $\mathbb{Z}[\beta]$ -linear combination of GQ_λ ’s, so the set $\{GQ_\lambda : \lambda \text{ is a strict partition}\}$ is a $\mathbb{Z}[\beta]$ -basis for a subring of $\mathbb{Z}[\beta][[x_1, x_2, \dots]]$.*

Proof. Theorem 7.8 implies that $GQ_\mu GQ_\nu = \sum_{\ell(\lambda) \leq n} b_{\mu\nu}^\lambda \beta^{|\lambda|-|\mu|-|\nu|} GQ_\lambda$ when $n = \ell(\mu) + \ell(\nu)$. This must be a finite sum since the set $\{GQ_\lambda(x_1, \dots, x_n) : \ell(\lambda) \leq n\}$ is a $\mathbb{Z}[\beta]$ -basis for a subring of $\mathbb{Z}[\beta][x_1, \dots, x_n]$ by [8, Prop. 3.2]. \square

For strict partitions $\mu \subseteq \lambda$ we may likewise define

$$(7.6) \quad JP_{\lambda/\mu} := \omega(GP_{\lambda/\mu}) \quad \text{and} \quad JQ_{\lambda/\mu} := \omega(GQ_{\lambda/\mu}),$$

setting $JP_\lambda := JP_{\lambda/\emptyset}$ and $JQ_\lambda := JQ_{\lambda/\emptyset}$. Interpreting these as combinatorial generating functions is easier than in the dual case. By [11, Cor. 6.6] we have

$$(7.7) \quad JP_{\lambda/\mu} = GP_{\lambda/\mu}\left(\frac{x}{1-\beta x}\right) \quad \text{and} \quad JQ_{\lambda/\mu} = GQ_{\lambda/\mu}\left(\frac{x}{1-\beta x}\right)$$

where $f\left(\frac{x}{1-\beta x}\right)$ denotes the power series obtained from $f \in \mathbb{Z}[\beta][[x_1, x_2, \dots]]$ by substituting $x_i \mapsto \frac{x_i}{1-\beta x_i} = x_i + \beta x_i^2 + \beta^2 x_i^3 + \dots$ for all i .

Using (7.7) it is straightforward to turn the formulas (5.3) into expressions for $JP_{\lambda/\mu}$ and $JQ_{\lambda/\mu}$ as generating functions $\sum_T \beta^{|T|-|\lambda/\mu|} x^T$ for *semistandard weak*

set-valued shifted tableau⁽³⁾ of shape λ/μ , with primed numbers excluded from the diagonal in the JP case. See [6, §3], for example, where the power series denoted K_λ is the same as $JP_\lambda|_{\beta=1}$. Since ω is linear, if we define

$$(7.8) \quad JP_{\lambda//\mu} := \omega(GP_{\lambda//\mu}) \quad \text{and} \quad JQ_{\lambda//\mu} := \omega(GQ_{\lambda//\mu})$$

then it also holds that

$$(7.9) \quad JP_{\lambda//\mu} = GP_{\lambda//\mu}\left(\frac{x}{1-\beta x}\right) \quad \text{and} \quad JQ_{\lambda//\mu} = GQ_{\lambda//\mu}\left(\frac{x}{1-\beta x}\right)$$

and it follows from (5.5) that

$$(7.10) \quad \begin{aligned} JP_\lambda(x, y) &= \sum_{\nu} JP_\nu(x)JP_{\lambda//\nu}(y), \\ JQ_\lambda(x, y) &= \sum_{\nu} JQ_\nu(x)JQ_{\lambda//\nu}(y), \end{aligned}$$

where the sums are over all strict partitions ν .

One can write down Cauchy identities relating each pair of GP/jq , GQ/jp , JP/gq , JQ/gp , JP/jq , and JQ/jp functions. These provide a shifted analogue of [24, Cor. 6.3]. Recall that $\bar{x}_i := \frac{-x_i}{1+\beta x_i}$ and $\Delta(x, y) := \prod_{i,j \geq 1} \frac{1-\bar{x}_i y_j}{1-x_i y_j}$.

COROLLARY 7.10. *Let μ and ν be strict partitions. Then:*

- (a) $\Delta(x, -y) \sum_{\lambda} GP_{\lambda//\mu}(x)jq_{\lambda/\nu}(y) = \sum_{\kappa} GP_{\nu//\kappa}(x)jq_{\mu/\kappa}(y);$
- (b) $\Delta(x, -y) \sum_{\lambda} GQ_{\lambda//\mu}(x)jp_{\lambda/\nu}(y) = \sum_{\kappa} GQ_{\nu//\kappa}(x)jp_{\mu/\kappa}(y);$
- (c) $\Delta(-x, y) \sum_{\lambda} JP_{\lambda//\mu}(x)gq_{\lambda/\nu}(y) = \sum_{\kappa} JP_{\nu//\kappa}(x)gq_{\mu/\kappa}(y);$
- (d) $\Delta(-x, y) \sum_{\lambda} JQ_{\lambda//\mu}(x)gp_{\lambda/\nu}(y) = \sum_{\kappa} JQ_{\nu//\kappa}(x)gp_{\mu/\kappa}(y);$
- (e) $\sum_{\lambda} JP_{\lambda//\mu}(x)jq_{\lambda/\nu}(y) = \Delta(-x, -y) \sum_{\kappa} JP_{\nu//\kappa}(x)jq_{\mu/\kappa}(y);$
- (f) $\sum_{\lambda} JQ_{\lambda//\mu}(x)jp_{\lambda/\nu}(y) = \Delta(-x, -y) \sum_{\kappa} JQ_{\nu//\kappa}(x)jp_{\mu/\kappa}(y).$

As usual the sums are over all strict partitions λ and κ .

Proof. As ω interchanges the elementary and complete symmetric functions $e_n := s_{1^n}$ and $h_n := s_{(n)}$, it follows that ω (extended to an automorphism of the ring of symmetric functions in the x_i variables over $\mathbb{Z}[t]$) also interchanges $E(t; x) := \sum_{n \geq 0} e_n t^n = \prod_{i \geq 1} (1 + x_i t)$ and $H(t; x) := \sum_{n \geq 0} h_n t^n = \prod_{i \geq 1} \frac{1}{1 - x_i t}$. The desired identities are straightforward to derive from Theorem 6.9 using this observation with (7.2) and (7.9), since one has

$$\Delta(x, y) = \prod_{i \geq 1} E(-\bar{x}_i; y)H(x_i; y) \quad \text{and} \quad -\bar{x}_i|_{x_i \mapsto \frac{x_i}{1-\beta x_i}} = \frac{\frac{x_i}{1-\beta x_i}}{1 + \beta \frac{x_i}{1-\beta x_i}} = x_i.$$

For example, substituting $x_i \mapsto \frac{x_i}{1-\beta x_i}$ and then applying the version of ω which acts only on symmetric functions in the y -variables transforms Theorem 6.9 to

$$\sum_{\lambda} JP_{\lambda//\mu}(x)jq_{\lambda/\nu}(y) = \prod_{i \geq 1} H(x_i; y)E\left(\frac{x_i}{1-\beta x_i}; y\right) \sum_{\kappa} JP_{\nu//\kappa}(x)jq_{\mu/\kappa}(y).$$

This implies (e) as $\prod_{i \geq 1} H(x_i; y)E\left(\frac{x_i}{1-\beta x_i}; y\right) = \prod_{i,j \geq 1} \frac{1 + \frac{x_i y_j}{1-\beta x_i}}{1 - x_i y_j} = \Delta(-x, -y)$. The other parts are derived in a similar way. \square

Acknowledgements. We thank Joel Lewis for several helpful discussions.

⁽³⁾A *semistandard weak set-valued shifted tableau* of shape λ/μ is defined in the same way as a semistandard set-valued shifted tableau, except the entries of such a tableau are finite nonempty multisets of positive half-integers.

REFERENCES

- [1] F. Barekat and S. van Willigenburg, *Composition of transpositions and equality of ribbon Schur Q -functions*, Electron. J. Comb. **16** (2009), no. 1, article no. R110.
- [2] A. S. Buch and V. Ravikumar, *Pieri rules for the K -theory of cominuscule Grassmannians*, J. fur Reine Angew. Math. **2012** (2012), no. 668, 109–132.
- [3] E. Clifford, H. Thomas, and A. Yong, *K -theoretic Schubert calculus for $OG(n, 2n+1)$ and jeu de taquin for shifted increasing tableaux*, J. fur Reine Angew. Math. **2014** (2014), no. 690, 51–63.
- [4] E. A. DeWitt, *Identities relating Schur s -functions and Q -functions*, Ph.D. thesis, University of Michigan, 2012.
- [5] M. Gillespie and K. Salois, *Inequality of a class of near-ribbon skew Schur Q -functions*, 2021, <https://arxiv.org/abs/2107.14212>.
- [6] Z. Hamaker, A. Keilthy, R. Patrias, L. Webster, Y. Zhang, and S. Zhou, *Shifted Hecke insertion and the K -theory of $OG(n, 2n+1)$* , J. Comb. Theory Ser. A **151** (2017), 207–240.
- [7] T. Hudson, T. Ikeda, T. Matsumura, and H. Naruse, *Double Grothendieck polynomials for symplectic and odd orthogonal Grassmannians*, J. Algebra **546** (2020), 294–314.
- [8] T. Ikeda and H. Naruse, *K -theoretic analogues of factorial Schur P - and Q -functions*, Adv. Math. **243** (2013), 22–66.
- [9] V. N. Ivanov, *Interpolation analogues of Schur Q -functions*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **307** (2004), no. Teor. Predst. Din. Sist. Komb. i Algoritm. Metody. 10, 99–119, 281–282.
- [10] T. Lam and P. Pylyavskyy, *Combinatorial Hopf Algebras and K -Homology of Grassmannians*, Int. Math. Res. Not. **2007** (2007), article no. rnm125.
- [11] J. B. Lewis and E. Marberg, *Enriched set-valued P -partitions and shifted stable Grothendieck polynomials*, Math. Z. (2021), 1929–1972.
- [12] I. G. Macdonald, *Symmetric functions and Hall polynomials*, 2 ed., Oxford university press, New York, 1999.
- [13] E. Marberg, *Shifted insertion algorithms for primed words*, 2021, <https://arxiv.org/abs/2104.11437>.
- [14] E. Marberg and B. Pawlowski, *K -theory formulas for orthogonal and symplectic orbit closures*, Adv. Math. **372** (2020), article no. 107299 (42 pages).
- [15] ———, *On some properties of symplectic Grothendieck polynomials*, J. Pure Appl. Algebra **225** (2021), no. 1, article no. 106463 (22 pages).
- [16] M. Nakagawa and H. Naruse, *Universal factorial Schur P, Q -functions and their duals*, 2018, <https://arxiv.org/abs/1812.03328>.
- [17] ———, *Generating functions for the universal factorial Hall-Littlewood P - and Q -functions*, 2021, <https://arxiv.org/abs/1705.04791v3>.
- [18] H. Naruse, *Elementary proof and application of the generating functions for generalized Hall-Littlewood functions*, J. Algebra **516** (2018), 197–209.
- [19] Soichi Okada, *A generalization of Schur's P - and Q -functions*, Sémin. Lothar. Combin. **81** (2020), article no. B81k (50 pages).
- [20] Oliver Pechenik and Alexander Yong, *Genomic tableaux*, J. Algebraic Combin. **45** (2017), no. 3, 649–685.
- [21] H. Salmasian, *Equality of Schur's Q -functions and Their Skew Analogues*, Ann. Comb. **3** (2008), no. 12, 325–346.
- [22] J. R. Stembridge, *Shifted tableaux and the projective representations of symmetric groups*, Adv. Math. **74** (1989), no. 1, 87–134.
- [23] ———, *Enriched P -partitions*, Trans. Amer. Math. Soc. **349** (1997), no. 2, 763–788.
- [24] D. Yeliussizov, *Symmetric Grothendieck polynomials, skew Cauchy identities, and dual filtered Young graphs*, J. Comb. Theory Ser. A **161** (2019), 453–485.

YU-CHENG CHIU, Department of Mathematics, ETH Zürich, Zürich, Switzerland
E-mail : chiuyu@student.ethz.ch

ERIC MARBERG, Department of Mathematics, HKUST, Clear Water Bay, Hong Kong
E-mail : emarberg@ust.hk